

# Microsporidia Evolved from Ancestral Sexual Fungi

Soo Chan Lee,<sup>1</sup> Nicolas Corradi,<sup>2</sup> Edmond J. Byrnes III,<sup>1</sup> Santiago Torres-Martinez,<sup>3</sup> Fred S. Dietrich,<sup>1,4</sup> Patrick J. Keeling,<sup>2</sup> and Joseph Heitman<sup>1,\*</sup>

<sup>1</sup>Department of Molecular Genetics and Microbiology  
Duke University Medical Center  
Durham, NC 27710  
USA

<sup>2</sup>Canadian Institute for Advanced Research  
Department of Botany  
University of British Columbia  
3529-6270 University Boulevard  
Vancouver, BC V6T1Z4  
Canada

<sup>3</sup>Departamento de Genética y Microbiología  
Facultad de Biología  
Universidad de Murcia  
30071 Murcia  
Spain

<sup>4</sup>Institute for Genome Sciences and Policy  
Duke University  
Durham, NC 27710  
USA

## Summary

Microsporidia are obligate, intracellular eukaryotic pathogens that infect animal cells, including humans [1]. Previous studies suggested microsporidia share a common ancestor with fungi [2–7]. However, the exact nature of this phylogenetic relationship is unclear because of unusual features of microsporidial genomes, which are compact with fewer and highly divergent genes [8]. As a consequence, it is unclear whether microsporidia evolved from a specific fungal lineage, or whether microsporidia are a sister group to all fungi. Here, we present evidence addressing this controversial question that is independent of sequence-based phylogenetic reconstruction, but rather based on genome structure. In the zygomycete basal fungal lineage, the *sex* locus is a syntenic gene cluster governing sexual reproduction in which a high mobility group (HMG) transcription-factor gene is flanked by triose-phosphate transporter (TPT) and RNA helicase genes [9]. Strikingly, microsporidial genomes harbor a *sex*-related locus with the same genes in the same order. Genome-wide synteny analysis reveals multiple other loci conserved between microsporidia and zygomycetes to the exclusion of all other fungal lineages with sequenced genomes. These findings support the hypothesis that microsporidia are true fungi that descended from a zygomycete ancestor and suggest microsporidia may have an extant sexual cycle.

## Results and Discussion

Microsporidia are unusual eukaryotic obligate intracellular pathogens that infect a wide variety of animals and cell types

[1]. The National Institute of Allergy and Infectious Diseases (NIAID) of the NIH includes microsporidia on the list of priority pathogens for biodefense research and the Environmental Protection Agency has also issued a caution concerning these waterborne pathogens. Microsporidia occur globally and infect the human gastrointestinal tract as a common cause of self-limited diarrheal disease in immunocompetent individuals or more chronic disease in immunocompromised hosts. Infections of other tissues, including the eye, the lung and the brain, also commonly occur [10]. Beyond humans, a variety of microsporidian species infect a wide range of animals, especially insects and fish, and in some cases are used for pest control [11, 12]. Infection by microsporidia involves an unusual mechanism in which eversion of the polar tube produces a syringe-like structure that pierces the target cell through which the infectious contents of the spore are rapidly injected into the host [13].

Within microsporidia, over 1,200 species in 150 genera have been described [1]; however, the phylogenetic placement of the group as a whole has been long debated. On the basis of an apparent absence of mitochondria, they were hypothesized to represent an ancient eukaryotic lineage [14, 15]. However, they are now known to harbor a mitochondrial remnant, the mitosome [16–19]. Phylogenetic studies also now argue against an early origin for microsporidia, suggesting instead that they are either derived from fungi or a sister group to the fungi [4, 6, 7, 20]. However, because the microsporidian genomes that have been studied encode few genes (<2,000) and the sequences of these genes are highly divergent [15, 21], molecular-phylogenetic reconstructions are difficult and notoriously unreliable [6, 8]. Accordingly, for correct definition of the relationship between microsporidia and fungi, other kinds of molecular characters may prove more useful than reconstructing phylogenies from divergent gene sequences. In this study we have examined a different kind of data that is not inherently affected by the accelerated rates of substitution common to microsporidian gene sequences. Specifically, we have examined genome-wide synteny and from this data demonstrate that: (1) microsporidia are true fungi; (2) they are specifically related to zygomycetes; and (3) they harbor genetic regulatory components that could function in sex determination and sexual reproduction.

Fungi undergo diverse, genetically programmed sexual development that is governed by the mating type (*MAT*) or sex locus [22, 23]. *MAT* loci in some fungi encode homeodomain proteins, and it has been proposed that microsporidian homeodomain genes may be the *MAT/sex* locus in *Encephalitozoon cuniculi* [24]. However, these homeodomain gene pairs differ from fungal *MAT* loci in several ways: They are colinear or convergent, rather than divergently transcribed, of the same functional class (HD2/HD2) or linked but not adjacent, and in no case is there synteny of flanking genes with known fungal *MAT* loci [25] (Figure S1 available online). There are reports suggesting that some microsporidian species may have extant sexual cycles on the sole basis of morphological observations and inferences [26, 27]. However, there is no molecular or genetic evidence suggesting that microsporidia undergo sexual reproduction, and there is limited genomic evidence (beyond

\*Correspondence: [heitm001@duke.edu](mailto:heitm001@duke.edu)




Figure 1. The sex Locus Is Conserved between Zygomycetes and Microsporidia

(A) Three zygomycetes and three microsporidia harbor a syntenic sex locus encoding a triose-phosphate transporter (TPT), HMG domain protein, and an RNA helicase. In zygomycetes, the orientation of *sexP* of *P. blakesleeaanus* is opposite to the other HMG protein genes and the *P. blakesleeaanus* (+) allele contains a repetitive element. The TPT gene of *R. oryzae* is inverted with respect to the *P. blakesleeaanus* and *M. circinelloides* sex alleles and an additional gene encoding a BTB-Ankyrin-RCC1 repeat protein is present. Microsporidia also have an additional hypothetical protein gene in the extended sex locus, and the RNA helicase and TPT genes are inverted with respect to the zygomycetes. Notably, in *A. locustae*, the RNA helicase gene lies at a different genomic location, and in *E. cuniculi* the HMG domain protein gene is flanked by a gene with more limited identity to HMG domain proteins.

(B) Dot plot analysis of the sex locus alleles of *M. circinelloides* and *P. blakesleeaanus* illustrates (+) and (-) specific DNA sequences at the sex locus. The x axis is the (-) and the y axis the (+) sex alleles. Sequences shown here are 7103 bp and 6511 bp for (+) and (-) strains of *M. circinelloides* and 9971 bp and 7631 bp for (+) and (-) strains of *P. blakesleeaanus*. The unique sequences at the sex locus span 1541 bp and 1463 bp for (+) and (-) strains of *M. circinelloides* and 5830 bp and 3494 bp for (+) and (-) strains of *P. blakesleeaanus*. The dot plot was performed with a 17 bp window with no mismatches allowed. "HP" stands for hypothetical protein.

both species, the TPT and RNA helicase genes are highly conserved between the (+) and (-) alleles, whereas the HMG genes (*sexP* and *sexM*) and their promoters have diverged (Figure 1B). The promoter regions of the TPT and RNA

helicase genes of *P. blakesleeaanus* are conserved in both mating types (Figure 1B), whereas in *M. circinelloides*, the RNA helicase gene promoter is conserved but the TPT promoter is within the sex locus and diverged (Figure 1B). As a consequence, the TPT genes in the (+) and (-) mating types of *M. circinelloides* might be differentially regulated during sexual development. The *R. oryzae* (-) sex locus has not yet been characterized.

that noted above for homeodomain gene homologs) for conserved machinery involved in sex in other organisms [28]. Thus, it has been unclear whether microsporidia undergo sexual reproduction and, if so, how it is genetically controlled. A recent study illustrated that the sex locus of the zygomycete fungi, *Phycomyces blakesleeaanus* and *Rhizopus oryzae*, represents a different paradigm compared to that of ascomycetes and basidiomycetes. In these species, the sex locus encodes a high-mobility group (HMG) domain protein flanked by genes encoding a triose-phosphate transporter (TPT) and an RNA helicase [9], forming a microsyntenic genomic locus in zygomycetous fungi (Figure 1A). Plus (+) and minus (-) mating-type strains of *P. blakesleeaanus* encode two divergent HMG proteins: *SexP* and *SexM*, respectively (Figure 1A). Searching the genome of the zygomycete, *Mucor circinelloides* (US Department of Energy Joint Genome Institute *Mucor* genome project) also revealed a *sexM* homolog in the (-) strain, and this *sexM* gene is also flanked by TPT and RNA helicase genes (Figure 1A). The corresponding sex locus (+) allele was isolated by PCR amplification and found to contain a *sexP* gene homolog between TPT and RNA helicase genes (Figure 1A). Two different (+) strains contained *sexP* and not *sexM*, whereas four different (-) strains harbored only *sexM*, consistent with mating-type tests (data not shown). In both mating types of *M. circinelloides*, the sex locus alleles are syntenic with flanking TPT and RNA helicase genes, as in *P. blakesleeaanus* (Figure 1A). In

comparison of gene order upstream and downstream of the *sexM* (*M. circinelloides* and *P. blakesleeaanus*) or *sexP* (*R. oryzae*) genes revealed that only the TPT, HMG, and RNA helicase genes are conserved and syntenic among the three zygomycetes (Figure S2). More importantly, the synteny of these three genes is not conserved in any other sequenced fungal genome, including representative species of the Ascomycota, Basidiomycota, and Chytridiomycota, where the TPT, HMG, and RNA helicase genes are unlinked and randomly distributed in the genomes (Figure S3). Thus, the extended sex locus (TPT, HMG [*SexP*/*SexM*], and RNA helicase) is highly specific to the Zygomycota and possibly to the Mucorales, the subphylum to which these species all belong.

Remarkably, given the lack of conservation of the extended sex locus in most fungi, we found that a syntenic gene cluster related to the sex locus is conserved in the genomes of three distantly related microsporidia (Figure 1A). In *Encephalitozoon cuniculi* and *Enterocytozoon bieneusi* the gene-order




Figure 2. Examples of Conserved Synteny and Relaxed Synteny Found in Zygomycetes and Microsporidia

(A) Conserved synteny among *P. blakesleeenanus*, *R. oryzae*, and the microsporidia. Syntenic 60S (L21) and 40S (S9) protein genes are conserved throughout fungi (see Figure S6). Another synteny of ribosomal protein and kinase genes is observed in two zygomycete genomes and two microsporidian genomes (*E. cuniculi* and *E. bienewisi*).

(B) Conserved synteny between *P. blakesleeenanus* and the three microsporidia.

(C) Conserved synteny between *R. oryzae* and the three microsporidia. White arrows indicate additional nonsyntenic genes. With the exception of two ribosomal proteins (L21 and S9), none of the syntenic gene clusters shown in panels (A), (B), and (C) were conserved in other ascomycete (*A. nidulans*), basidiomycete (*C. neoformans*), or chytridiomycete (*B. dendrobatidis*) fungi. The genomes of *P. blakesleeenanus* and *R. oryzae* were compared to the *E. cuniculi* genome with BLASTp. Homologs in *E. bienewisi* and *A. locustae* were identified with tBLASTn.

conservation extends in both directions (with an additional ORF between TPT and the putative sex locus and inversions of flanking genes), whereas in *Antonospora locustae* the RNA helicase is no longer linked (Figure 1A). The gene order around the sex-related locus of both *E. cuniculi* and *A. locustae* was verified by PCR and sequencing. Interestingly, in both *R. oryzae* and the microsporidians, the TPT gene is inverted relative to other zygomycetes, and an additional gene has also been acquired between it and the sex locus (Figure 1A). It is possible that independent inversion events introduced these new genes into the extended sex locus. Structural differences of the sex locus observed in zygomycetes and microsporidia suggest alternative hypotheses for the early steps in the evolution of sex determination and sex chromosomes [9] (Figure S4).

To investigate whether the high degree of genome structure conservation between microsporidia and zygomycetes is restricted to the sex related locus, we also performed whole-genome comparisons between fungi and microsporidia. Interestingly, in a relaxed synteny analysis that allows for local changes in gene order or orientation to be accommodated (see Experimental Procedures and Figure S5) but still requires

a higher level genomic architecture be apparent, microsporidia and the zygomycetes were found to contain a number of conserved pairs or clusters of genes that were not conserved in any other fungal genome (BLAST cut off  $1 \times e^{-30}$ ) (Figure 2). The number of syntenic gene loci (33 gene pairs or clusters) shared between *R. oryzae* and *E. cuniculi* is significant compared to randomized gene orders ( $17.4 \pm 4.4$  pairs) (BLAST cut off  $1 \times e^{-05}$ ) (Figure S5 and Table S1). On the other hand, by the same analysis, the number of genomic regions that exhibit relaxed synteny and that are shared between *E. cuniculi* and other fungi, including *Saccharomyces cerevisiae*, *Ashbya gossypii*, *Neurospora crassa*, *Schizosaccharomyces pombe*, and *Cryptococcus neoformans*, is not significantly different in comparison to the randomized gene orders (Figure S6). The only gene pair that was conserved in a broader range of fungi, including the three microsporidia, was *RPL21* and *RPS9*, which is intriguingly conserved across all fungi (Figure S7) but not in species outside the fungal kingdom, suggesting an ancient functional role for this gene pair within the fungi. The shared, genome-wide preservation of synteny is significant because it is easy to imagine that synteny around the sex locus could be conserved as a result of functional

constraints, and so multiple independent losses of this synteny in other fungal lineages could reflect functional shifts in mating-type determination. The striking degree of overall conservation specifically between microsporidia and zygomycetes is not easily explained unless microsporidia and zygomycetes share a recent common ancestor relative to other fungi.

To explain the genome data in any other way (e.g., if microsporidia are the sister group to all fungi), one would have to postulate that the genomes of certain members of both microsporidia and zygomycetes were somehow special in having independently retained a high proportion of the gene order that was present in the ancestor of microsporidia and all fungi and that was lost multiple times independently in other fungal lineages. Indeed, it has been demonstrated that gene order evolves slowly in microsporidian genomes [21, 29]; however, the extended conservation in synteny that we document is not consistent with the idea that the same is true of zygomycetes. If that were true, multiple syntenic regions should be shared by all zygomycete and microsporidian genomes, but aside from the *sex* and *RPL21/RPS9* loci, this is not the case (Figure 2). Instead, gene order is typically conserved in all three microsporidia, but only one or two of the zygomycetes. In addition, the overall level of synteny between different zygomycetes does not suggest the same level of conservation found in microsporidia. This is all consistent with slow rearrangements in microsporidian genomes [21, 29], but not consistent with a slow rate of rearrangements in zygomycete genomes. There is accordingly no evidence that the gene order retained by zygomycetes and microsporidia is ancestral to all fungi.

Support for either explanation will be bolstered significantly by the additional sampling of basal fungal genomes. If other species of chytrids or “zygomycetes” not specifically related to Mucorales are subsequently found to share a significant fraction of the synteny described here, then it will suggest that the synteny may be derived from the ancestor of all fungi. If, on the other hand, no conservation is detected in these genomes, it further supports the relationship between microsporidia and Mucorales. Indeed, it is also possible that other subgroups of zygomycetes may share an even closer relationship, such as the Entomophthorales, which were proposed to be closely related to microsporidia on the basis of molecular phylogenies [4]. Whatever the outcome, the availability of additional zygomycete, chytridiomycete, and microsporidian genome sequences that are currently in progress will allow repeated tests of their relationships. However, on the basis of the genomes available now, the close evolutionary relationship between microsporidia and zygomycetes is the most parsimonious explanation of the data.

Sexual reproduction in microsporidia is poorly studied at best and completely unknown in most species, so the identification of a putative sex-determining locus in microsporidia has functional implications deserving of further study. Analysis of other isolates of the species described here may reveal the existence of divergent HMG proteins, as in the (+) and (–) mating types of zygomycetes. However, it is also possible they are homothallic. The HMG domain proteins of *E. cuniculi* and *A. locustae* each contain two HMG domains, whereas SexP and SexM of the zygomycetes contain a single HMG domain. The two microsporidian HMG domains are linked by 24 or 27 amino acid spacers respectively (Figure S8), suggesting the two domains could interact with DNA independently [30]. If these represent functional correlates of SexP and SexM, and each domain in a single protein confers both SexP and SexM

activity, the microsporidia might be homothallic and self-fertile. Maximum-likelihood analysis showed that both microsporidian HMG domains are more closely related to SexP than to SexM (Figure S8). In view of this, we propose three possible models for sexual development of microsporidia: (1) homothallism with a tandem HMG domain protein, (2) heterothallism with divergent HMG alleles that confer opposite mating types, as in zygomycetes, and (3) loss of sex [31], in which case the sex-related locus is an evolutionary vestige. Opportunities for sexual reproduction in microsporidia could occur during coinfection or involve self-fertile, homothallic mating, or interactions between nuclei in species known to be dikaryotic. Of note, the hypothesis of extant sexual cycles is further supported by the finding that a variety of meiotic gene homologs are present in all three of the microsporidian genomes analyzed here (Table S2), some of which were noted for *E. cuniculi* in a recent report by other investigators [28].

The high degree of synteny shared between microsporidian genomes and zygomycete genomes, including the sex-related locus, provides a compelling new line of evidence to resolve the enigmatic origin of these unusual eukaryotic intracellular pathogens. These data demonstrate that microsporidia are true fungi that share a close common ancestor with zygomycetes, and perhaps specifically with the Mucorales. The data also suggest a new avenue to examine the difficult problem of sex in microsporidia. At present, there is no direct, functional link between the sex-locus-related gene cluster in the microsporidia and mating or meiosis. No sexual cycle has been unambiguously defined for the three microsporidian species considered here and inferences about whether other species even have a sexual cycle are limited. However, the finding that they share a surprisingly well-conserved syntenic sex locus that governs mating in zygomycetous fungi leads to the reasonable hypothesis that it may also control sex in microsporidia. Future studies should address whether coinfection with genetically distinct isolates leads to the production of recombinant progeny, analysis of if and when the genes encoded by the sex-locus-related gene cluster are expressed, and analysis of whether other genes are present that might be inferred to play roles in sexual reproduction (such as genes encoding enzymes involved in production or sensing of the zygomycete trisporic acid pheromone or sex and meiosis-related functions) [28]. Our findings that microsporidia evolved from a basal fungal lineage further invites speculation about whether the microsporidian polar-tube infection structure is related to the sporangiophores of zygomycetous fungi, with implications for its functional roles in sensing extracellular signals and interactions with the host.

#### Experimental Procedures

To identify the sex locus of *M. circinelloides*, we performed tBLASTn analysis with the amino acid sequence of *P. blakesleeenae* SexM against the 8× genome sequence of R7B, a (–) strain of *M. circinelloides* (US Department of Energy Joint Genome Institute *Mucor* genome project, <http://mucorgen.un.es>). The DNA sequence was annotated with FGENESH or Orf finder at NCBI. (–) strains (R7B, ATCC1216b, CBS277.49, and NRRL1443) were used to confirm the (–) sex locus. We used genomic DNAs of (+) strains (NRRL3631 and ATCC1216a) to amplify, identify, and sequence the (+) sex allele.

Comparison of gene order across genomes was carried out starting with the annotated protein set for *E. cuniculi* [15], *R. oryzae* (Broad Institute Database for *R. oryzae*), and *P. blakesleeenae* (US Department of Energy Joint Genome Institute *P. blakesleeenae* genome project). Blast libraries were created with the NCBI blast database formatting tool formatdb [32]. The *P. blakesleeenae* and *R. oryzae* genome protein sets were compared

against the *E. cuniculi* genome protein set, and we compared each protein set to their own genomic DNA sequence to ascertain the order of genes within the genome. Genes were classified as homologs between these species in cases in which BLASTp E-values were less than  $1.0 \times e^{-30}$ . Areas sharing relaxed synteny were defined in which two or more gene homologs ( $1.0 \times e^{-10}$ ) were clustered and in the same order but included cases in which orientation differed or in which one or a few genes had been interposed (see Figure 2 and Figure S5 and legend for examples and details).

#### Accession Numbers

*Mucor circinelloides* sex locus sequences have been deposited in GenBank under accession FJ009106 and FJ009107. Microsporidian sex-related locus sequences have been deposited in GenBank under accession FJ008719 and FJ008720.

#### Supplemental Data

Supplemental Data include two tables and eight figures and can be found with this article online at [http://www.current-biology.com/supplemental/S0960-9822\(08\)01256-6](http://www.current-biology.com/supplemental/S0960-9822(08)01256-6).

#### Acknowledgments

We acknowledge access to the *Mucor circinelloides* and *Phycomyces blakesleeenanus* genome data established by the US Department of Energy Joint Genome Institute and to the *Rhizopus oryzae* genome data generated by the Broad Institute/Fungal Genome Initiative. We thank Donna Akiyoshi for the access to the *E. bienersi* genome database. The *Enterocytozoon bienersi* genome sequence survey was supported by National Institutes of Health grants R21 AI52792 and R21 AI064118. We thank Hilary Morrison and acknowledge the Josephine Bay Paul Center for Comparative Molecular Biology and Evolution for the use of data included in the *Antonosporea locustae* Genome Project funded by NSF award number 0135272. We thank Rytas Vilgalys, Andrii Gryganskiy, Tim James, Alex Idnurm, Raphael Valdivia, Banu Metin, Charles Li, and Naomi Fast for discussions or comments on the manuscript. This work was supported by National Institutes of Health grant AI50113 to J.H. and by Canadian Institutes for Health Research grant MOP-84265 to P.J.K. P.J.K. is a Fellow of the Canadian Institute for Advanced Research and a Senior Scholar of the Michael Smith Foundation for Health Research.

Received: July 18, 2008

Revised: August 29, 2008

Accepted: September 1, 2008

Published online: October 30, 2008

#### References

- Keeling, P.J., and Fast, N.M. (2002). Microsporidia: Biology and evolution of highly reduced intracellular parasites. *Annu. Rev. Microbiol.* 56, 93–116.
- Edlind, T.D., Li, J., Visvesvara, G.S., Vodkin, M.H., McLaughlin, G.L., and Katiyar, S.K. (1996). Phylogenetic analysis of beta-tubulin sequences from amitochondrial protozoa. *Mol. Phylogenet. Evol.* 5, 359–367.
- Keeling, P.J., and Doolittle, W.F. (1996). Alpha-tubulin from early-diverging eukaryotic lineages and the evolution of the tubulin family. *Mol. Biol. Evol.* 13, 1297–1305.
- Keeling, P.J. (2003). Congruent evidence from alpha-tubulin and beta-tubulin gene phylogenies for a zygomycete origin of microsporidia. *Fungal Genet. Biol.* 38, 298–309.
- James, T.Y., Kauff, F., Schoch, C.L., Matheny, P.B., Hofstetter, V., Cox, C.J., Celio, G., Gueidan, C., Fraker, E., Miadlikowska, J., et al. (2006). Reconstructing the early evolution of Fungi using a six-gene phylogeny. *Nature* 443, 818–822.
- Hirt, R.P., Logsdon, J.M., Healy, B., Dorey, M.W., Doolittle, W.F., and Embley, T.M. (1999). Microsporidia are related to Fungi: Evidence from the largest subunit of RNA polymerase II and other proteins. *Proc. Natl. Acad. Sci. USA* 96, 580–585.
- Gill, E.E., and Fast, N.M. (2006). Assessing the microsporidia-fungi relationship: Combined phylogenetic analysis of eight genes. *Gene* 375, 103–109.
- Thomarat, F., Vivarès, C.P., and Gouy, M. (2004). Phylogenetic analysis of the complete genome sequence of *Encephalitozoon cuniculi* supports the fungal origin of microsporidia and reveals a high frequency of fast-evolving genes. *J. Mol. Evol.* 59, 780–791.
- Idnurm, A., Walton, F.J., Floyd, A., and Heitman, J. (2008). Identification of the sex genes in an early diverged fungus. *Nature* 451, 193–196.
- Didier, E.S., and Weiss, L.M. (2006). Microsporidiosis: Current status. *Curr. Opin. Infect. Dis.* 19, 485–492.
- Lom, J., and Nilsen, F. (2003). Fish microsporidia: Fine structural diversity and phylogeny. *Int. J. Parasitol.* 33, 107–127.
- Becnel, J.J., and Andreadis, T.G. (1999). Microsporidia in insects. In *The microsporidia and microsporidiosis*, M. Wittner, ed. (Washington, D.C.: ASM Press), pp. 1–6.
- Xu, Y., and Weiss, L.M. (2005). The microsporidian polar tube: A highly specialised invasion organelle. *Int. J. Parasitol.* 35, 941–953.
- Vossbrinck, C.R., Maddox, J.V., Friedman, S., Debrunner-Vossbrinck, B.A., and Woese, C.R. (1987). Ribosomal-RNA sequence suggests microsporidia are extremely ancient eukaryotes. *Nature* 326, 411–414.
- Katinka, M.D., Duprat, S., Cornillot, E., Metenier, G., Thomarat, F., Prensier, G., Barbe, V., Peyretailade, E., Brottier, P., Wincker, P., et al. (2001). Genome sequence and gene compaction of the eukaryote parasite *Encephalitozoon cuniculi*. *Nature* 414, 450–453.
- Tsaousis, A.D., Kunji, E.R.S., Goldberg, A.V., Lucocq, J.M., Hirt, R.P., and Embley, T.M. (2008). A novel route for ATP acquisition by the remnant mitochondria of *Encephalitozoon cuniculi*. *Nature* 453, 553–556.
- Williams, B.A.P., Hirt, R.P., Lucocq, J.M., and Embley, T.M. (2002). A mitochondrial remnant in the microsporidian *Trachipleistophora hominis*. *Nature* 418, 865–869.
- Goldberg, A.V., Molik, S., Tsaousis, A.D., Neumann, K., Kuhnke, G., Delbac, F., Vivares, C.P., Hirt, R.P., Lill, R., and Embley, T.M. (2008). Localization and functionality of microsporidian iron-sulphur cluster assembly proteins. *Nature* 452, 624–628.
- Williams, B.A.P., Cali, A.N.N., Takvorian, P.M., and Keeling, P.J. (2008). Distinct localization patterns of two putative mitochondrial proteins in the microsporidian *Encephalitozoon cuniculi*. *J. Eukaryot. Microbiol.* 55, 131–133.
- Keeling, P.J., Luker, M.A., and Palmer, J.D. (2000). Evidence from beta-tubulin phylogeny that microsporidia evolved from within the fungi. *Mol. Biol. Evol.* 17, 23–31.
- Slamovits, C.H., Fast, N.M., Law, J.S., and Keeling, P.J. (2004). Genome compaction and stability in microsporidian intracellular parasites. *Curr. Biol.* 14, 891–896.
- Dyer, P.S. (2008). Evolutionary biology: Genomic clues to original sex in fungi. *Curr. Biol.* 18, R207–R209.
- Fraser, J.A., and Heitman, J. (2003). Fungal mating-type loci. *Curr. Biol.* 13, R792–R795.
- Bürglin, T.R. (2003). The homeobox genes of *Encephalitozoon cuniculi* (Microsporidia) reveal a putative mating-type locus. *Dev. Genes Evol.* 213, 50–52.
- James, T.Y. (2007). Analysis of mating-type locus organization and synteny in mushroom fungi: Beyond model species. In *Sex in Fungi*, J. Heitman, J.W. Kronstad, J.W. Taylor, and L.A. Casselton, eds. (Washington, DC: ASM Press), pp. 317–331.
- Hazard, E.I., and Brookbank, J.W. (1984). Karyogamy and meiosis in an *Amblyospora* Sp (Microspora) in the mosquito *Culex salinarius*. *J. Invertebr. Pathol.* 44, 3–11.
- Sweeney, A.W., Graham, M.F., and Hazard, E.I. (1988). Life cycle of *Amblyospora dyxenooides* sp. nov. in the mosquito *Culex annulostris* and the copepod *Mesocyclops albianus*. *J. Invertebr. Pathol.* 51, 46–57.
- Malik, S.-B., Pightling, A.W., Stefaniak, L.M., Schurko, A.M., and Logsdon, J.M., Jr. (2008). An expanded inventory of conserved meiotic genes provides evidence for sex in *Trichomonas vaginalis*. *PLoS ONE* 3, e2879.
- Corradi, N., Akiyoshi, D.E., Morrison, H.G., Feng, X., Weiss, L.M., Tzipori, S., and Keeling, P.J. (2007). Patterns of genome evolution among the microsporidian parasites *Encephalitozoon cuniculi*, *Antonosporea locustae* and *Enterocytozoon bienersi*. *PLoS ONE* 2, e1277.
- Stott, K., Tang, G.S.F., Lee, K.-B., and Thomas, J.O. (2006). Structure of a complex of tandem HMG boxes and DNA. *J. Mol. Biol.* 360, 90–104.
- Ironside, J. (2007). Multiple losses of sex within a single genus of microsporidia. *BMC Evol. Biol.* 7, 48.
- Pearson, W.R., and Lipman, D.J. (1988). Improved tools for biological sequence comparison. *Proc. Natl. Acad. Sci. USA* 85, 2444–2448.