



Invited Review

The search for the missing link: A relic plastid in *Perkinsus*?

José A. Fernández Robledo^{a,*}, Elisabet Caler^b, Motomichi Matsuzaki^c, Patrick J. Keeling^d,
Dhanasekaran Shanmugam^e, David S. Roos^e, Gerardo R. Vasta^a

^a Department of Microbiology and Immunology, School of Medicine, University of Maryland, IMET, 701 East Pratt Street, Suite 236, Baltimore, MD 21202-3101, USA

^b J. Craig Venter Institute, 9704 Medical Center Drive, Rockville, MD 20850-3343, USA

^c Department of Biomedical Chemistry, Graduate School of Medicine, University of Tokyo, Japan

^d Department of Botany, Canadian Institute for Advanced Research, University of British Columbia, Vancouver, BC, Canada

^e Department of Biology and Penn Genome Frontiers Institute, University of Pennsylvania, Philadelphia, PA 10104, USA

ARTICLE INFO

Article history:

Received 30 June 2011

Received in revised form 27 July 2011

Accepted 28 July 2011

Available online 22 August 2011

Keywords:

Apicoplast

Chromalveolata hypothesis

Genome

Perkinsus

Plastid

Protozoan

Targeting

ABSTRACT

Perkinsus marinus (Phylum Perkinsozoa) is a protozoan parasite that has devastated natural and farmed oyster populations in the USA, significantly affecting the shellfish industry and the estuarine environment. The other two genera in the phylum, *Parvilucifera* and *Rastrimonas*, are parasites of microeukaryotes. The Perkinsozoa occupies a key position at the base of the dinoflagellate branch, close to its divergence from the Apicomplexa, a clade that includes parasitic protista, many harbouring a relic plastid. Thus, as a taxon that has also evolved toward parasitism, the Perkinsozoa has attracted the attention of biologists interested in the evolution of this organelle, both in its ultrastructure and the conservation, loss or transfer of its genes. A review of the recent literature reveals mounting evidence in support of the presence of a relic plastid in *P. marinus*, including the presence of multimembrane structures, characteristic metabolic pathways and proteins with a bipartite N-terminal extension. Further, these findings raise intriguing questions regarding the potential functions and unique adaptation of the putative plastid and/or plastid genes in the Perkinsozoa. In this review we analyse the above-mentioned evidence and evaluate the potential future directions and expected benefits of addressing such questions. Given the rapidly expanding molecular/genetic resources and methodological toolbox for *Perkinsus* spp., these organisms should complement the currently established models for investigating plastid evolution within the Chromalveolata.

© 2011 Published by Elsevier Ltd. on behalf of Australian Society for Parasitology Inc.

1. Introduction

1.1. Life cycle

Perkinsus spp. have a direct life cycle (Fig. 1): trophozoites proliferate intra- or extracellularly by palintomy (merogony or schizogony) giving rise to 4–32 (often 8–16) trophozoites, which are released upon rupture of the schizont cell wall (Perkins, 1996; Sunila et al., 2001). At the four-cell schizont stage, a functional diversification of the daughter cells has been proposed (Sunila et al., 2001). Whilst less frequent, binary fission has also been observed: trophozoite budding yields a mother cell surrounded by a thick wall and the daughter cell separated by a plasma membrane (Sunila et al., 2001). Sexual stages have been suggested (Perkins, 1996) and recent microsatellite analyses suggest that *Perkinsus marinus* utilises both sexual and asexual reproduction,

and that over the short term selection acts upon independent parasite lineages rather than upon individual loci in a cohesive, interbreeding population (Thompson et al., 2011). Although the overall propagation process has been described in substantial detail, various aspects of cell division and maturation at the subcellular level remain poorly understood, especially regarding the segregation and function of some organelles and macrostructures (e.g. the large vacuole containing the vacuoplast and its precursors during trophozoite proliferation and maturation) and the zoosporulation process.

1.2. Natural history and phylogenetic position

In the early 1950s, the causative agent of widespread mass mortalities of eastern oysters (*Crassostrea virginica*) on the shores of Texas (USA) was identified and named *Dermocystidium marinum*. This microorganism was later renamed *Perkinsus marinus* and closely related *Perkinsus* spp. that affect various mollusc species worldwide were described during later years (Villalba et al., 2004). The gradual expansion of the geographic distribution of

* Corresponding author. Tel.: +1 410 234 8849; fax: +1 410 234 8896.

E-mail address: jfernandez-robledo@som.umaryland.edu (J.A. Fernández Robledo).





Fig. 1. *Perkinsus* life cycle. (A) Trophozoites in the water column (1) are taken by the oyster during filter-feeding (2), enter the paleal cavity and are directed through gills and palps towards the mouth. Trophozoites may be released into the water (3) from live oysters together with the pseudofeces and upon death of the oyster, from the decaying infected tissues (Bushek et al., 2002). Once released into the water column, trophozoites may sporulate (4): trophozoites enlarge, develop a discharge tube and after multiple rounds of division, release hundreds of zoospores into the water column. Whether zoospores develop into trophozoites remains an open question. (B) Once in the paleal cavity or the digestive tract (5), trophozoites displaying surface ligands for the oyster galectin CvGal (Tasumi and Vasta, 2007) are recognised and phagocytosed by the hemocytes (6) that can translocate to the internal milieu (7) and eventually into the vascular system (8). Parasites remain inside phagosome-like vesicles where they remain viable and multiply. When hemocytes disintegrate (9, 10), the released trophozoites can either be phagocytosed by neighbouring hemocytes or multiply extracellularly in both the internal milieu (11) and arterial lumen (12). The infected circulating hemocytes migrate throughout the host tissues where they may lyse and release trophozoites, leading to systemic infection and eventually host death. (C) *In vitro* culture of *Perkinsus marinus* trophozoites. Under scanning electron microscopy the cultured *P. marinus* trophozoite surface appears smooth (a, b). In trophozoites undergoing schizogony, the shapes of the daughter cells become apparent on the exterior surface (stars, b, c).

P. marinus infections along the Atlantic coast of the USA has been associated with global warming and shellfish trade (Ford, 1996; Ford and Chintala, 2006; Ford and Smolowitz, 2007; Pecher et al., 2008), and it is currently under surveillance by the World

Organization for Animal Health (<http://www.oie.int/>). In addition to constituting a valuable shellfisheries resource, filter-feeding bivalves are important components of marine and estuarine environments because they play a critical role in maintaining water quality and ecosystem integrity. Since its discovery, *P. marinus* has been placed in various taxa including fungi, saprolegniales, apicomplexans and dinozoans (Mackin et al., 1950; Mackin and Ray, 1966; Levine, 1978; Goggin and Barker, 1993; Siddall et al., 1997; Kuvardina et al., 2002). More recently, the new phylum Perkinsozoa was established to include *Perkinsus* and two other genera, *Parvilucifera* and *Rastrimonas*, which are parasites of microeukaryotes. *Parvilucifera* is a parasite of the toxic dinoflagellate *Dinophysis* (Norén et al., 1999), and *Rastrimonas* infects the free-living cryptophyte *Chilomonas paramecium* (Brugerolle, 2002). The Perkinsozoa is considered to be the earliest group diverging from the lineage leading to dinoflagellates, branching close to the node shared by dinoflagellates and apicomplexans (Saldarriaga et al., 2003; Gile et al., 2006; Moore et al., 2008; Bachvaroff et al., 2011). Members of both dinoflagellates and apicomplexans possess plastids and recent analysis of newly identified photosynthetic members of the apicomplexan lineage have shown that these plastids evolved from a single common secondary endosymbiosis with a red alga (Moore et al., 2008; Janouskovec et al., 2010). This suggests that non-photosynthetic relatives of both lineages, including *Perkinsus*, evolved from photosynthetic ancestors, raising the possibility that these lineages retain cryptic organelles (Keeling, 2010). Similarly, expressed sequence tag (EST) evidence suggests that the non-photosynthetic early-diverging dinoflagellate, *Cryptocodinium cohnii*, may contain a reduced plastid (Sánchez-Puerta et al., 2007). Here, we analyse the current evidence in support of the presence of a relic plastid in *Perkinsus*, as well as the remaining open questions. We also evaluate the potential future directions and expected benefits of rigorously addressing such questions in a functional and evolutionary context within the Chromalveolata. The Chromalveolata is a eukaryote supergroup established to include those taxa that comprise organisms resulting from a single secondary endosymbiosis between a line descending from a bikont and a red alga (Cavalier-Smith, 1999; Keeling, 2010). It includes the Alveolata, a group of unicellular eukaryotes characterised by the presence of cortical alveoli (Cavalier-Smith, 1998).



2. The Chromalveolata hypothesis: the apicoplast, a relic plastid in apicomplexans

2.1. Origin and architecture of the apicoplast

The apicoplast was described over two decades ago in several apicomplexan parasites as a multimembrane vesicle (Kilejian, 1974;

Fig. 2. Plastid origin and evolution. (A) Under the Chromalveolata hypothesis (Cavalier-Smith, 1999; Keeling, 2010) eukaryotic plastids are DNA-containing organelles that have evolved via secondary endosymbiosis resulting from the initial engulfment of a free-living photosynthetic cyanobacterium by a heterotrophic protist, followed by its secondary engulfment by a heterotrophic eukaryotic host. The hallmark of this process is the presence of a three- to four-membrane organelle that has maintained a remnant of the original cyanobacterium genome and essential metabolic functions. (B) Both the Apicomplexa and Perkinsida are constituted by parasitic protista that have lost photosynthetic capabilities. (C) In addition to genes encoded by the plastid genome derived from the secondary endosymbiosis in apicomplexans and perkinsids, approximately 500 nuclear-encoded apicomplexan proteins are predicted to be targeted back to the plastid through a bipartite targeting sequence (as depicted in D); similarly, nuclear genes encoding putative plastid pre-proteins containing an N-terminal signal peptide have been identified in *Perkinsus*. (D) In apicomplexans, the nuclear-encoded plastid pre-proteins contain an N-terminal signal peptide (SP) which targets the protein to the secretory pathway, followed by a transit peptide (TP) which targets the protein to the plastid; whether this process also takes place in perkinsids remains an open question. N, nucleus. MP, mature peptide.

Gardner et al., 1991; Howe, 1992; Williamson et al., 1994; McFadden et al., 1996; Wilson et al., 1996; Köhler et al., 1997). Its ultrastructural hallmark is the presence of four intimately associated membranes, although three membranes have been reported in some instances (e.g. *Plasmodium* spp.), whilst in others the organelle is completely absent (e.g. *Cryptosporidium* spp.) (Hopkins et al., 1999; Zhu et al., 2000; Abrahamsen et al., 2004; Xu et al., 2004). The Chromalveolata hypothesis (Cavalier-Smith, 1999) (Fig. 2A,B) proposed that apicomplexan plastids resulted from a single secondary endosymbiosis of a red alga that also gave rise to plastids in dinoflagellates, heterokonts, haptophytes and cryptomonads (reviewed in Keeling, 2010). Whilst this hypothesis remains contentious for several of these lineages, there is now strong evidence that at least the apicomplexan and dinoflagellate plastids did arise from a common ancestor. Both of these organelles are unusual and have proved difficult to compare but the recent discovery of photosynthetic relatives of apicomplexans (Moore et al., 2008) and the characterisation of their plastid genomes and associated genes (Janouskovec et al., 2010) have provided multiple lines of evidence for their common ancestry. Both primary and secondary endosymbiotic events are followed by drastic reductions in the genome of the engulfed endosymbiont. For example, because a chloroplast encodes for only 5–10% of the genes present in the free-living cyanobacteria, it is estimated that between 800 to 2,000 genes from the original endosymbiont were transferred to the nucleus of the primary host (Martin and Herrmann, 1998; Martin et al., 2002), and in extant species, most of the proteins targeted in a plastid are nuclear encoded. The apicomplexan plastid retained a 35–40-kb circular extrachromosomal genetic element considered as a remnant of a secondary endosymbiotic event (Fig. 2C,D), and houses 33 tRNA (e.g. *Toxoplasma* *plDNA*), two head-to-head copies of rRNA genes, RNA polymerase genes, numerous housekeeping genes (mostly related to gene expression) and several open reading frames (ORFs) (Feagin and Parsons, 2007; Kissinger and Kuo, 2007). In dinoflagellates this reduction has been even more extreme: with only 13 genes, mostly relating to photosystems, retained on single-gene mini-circle chromosomes, they have the most reduced plastid genomes documented to date (Green, 2011) and at least some of the many missing genes are known to be located in the nucleus (Bhattacharya et al., 2004).

2.2. Metabolic relevance of the apicoplast

Key metabolic functions associated with the apicoplast include complete pathways for isoprenoid synthesis using the non-mevalonate pathway (MEP pathway), type II fatty acid synthesis (FAS), lipoic acid, abscisic acid (ABA), iron–Sulphur (Fe–S) cluster and haeme biosynthesis; in addition the apicoplast harbours certain steps of glycolysis and phospholipid synthesis (reviewed in Seeber and Soldati-Favre, 2010). The MEP and type II FAS pathways are absent in mammals and other enzymes (e.g. Porphobilinogen synthase (PBGs) in haeme biosynthesis) differ from the human counterpart; hence, the enzymes involved in these pathways have been recognised as promising drug targets (Fichera and Roos, 1997; McFadden and Roos, 1999; Soldati, 1999; Wiesner et al., 2008; Lim et al., 2009; Jaffe et al., 2011). Currently, on a quest for new drugs to treat diseases caused by apicomplexan parasites, additional metabolic pathways associated with the apicoplast as well as other aspects of basic plastid biology are being characterised as potential targets for intervention (Muller and Hemphill, 2011).

3. *Perkinsus marinus* is a key organism for understanding plastid evolution

The Perkinsozoa (*Perkinsus*, *Parvilucifera* and *Rastrimonas*) is positioned at the base of the dinoflagellate branch close to the

divergence from the Apicomplexa (Goggin and Barker, 1993; Reece et al., 1997; Siddall et al., 1997; Ellis et al., 1998; Norén et al., 1999; Leander and Keeling, 2003) and represents a key taxon for understanding adaptations and organelle evolution within the Chromalveolata. Indeed, structures resembling a plastid have been identified in *Rastrimonas subtilis*: “Two enigmatic bodies limited by three membranes surrounding a matrix denser at the periphery than in the center” (Brugerolle, 2002), and in *Parvilucifera infectans*: “an unidentified organelle surrounded by two membranes... with the occasional indication of a third. They appear to represent small mitochondria...” (Norén et al., 1999). Interestingly, when *Parvilucifera prorocentri* was recently described, no bona fide apicoplast homologue was identified (Leander and Hoppenrath, 2008). Similarly, no reference was made to any plastid or enigmatic bodies in the description of *Parvilucifera sinerae* (Figuroa et al., 2008). Unfortunately, the lack of sequenced genomes for these species has precluded their mining for evidence of a plastid. Nevertheless, increasing ultrastructural and molecular evidence discussed below from *Perkinsus* points towards the presence of a relic plastid, which together with the currently available resources and methodological tools (two fully defined culture media formulations (Gauthier et al., 1995; La Peyre and Faisal, 1997), a large number of *Perkinsus* isolates and species (<http://www.atcc.org>), a *P. marinus* sequenced genome, a transcriptome (Joseph et al., 2010) and a transfection system (Fernández-Robledo et al., 2008a)) make *Perkinsus* spp. a potentially unique model system to gain further insight into plastid evolution in this non-photosynthetic sister lineage to dinoflagellates.

3.1. A *Perkinsus* plastid genome?

The identification of three plastid-associated major biosynthetic pathways (MEP (Matsuzaki et al., 2006, 2008; Grauvogel et al., 2007; Joseph et al., 2010), type II FAS and Fe–S cluster (Stelter et al., 2007)) in *Perkinsus* suggests the presence of a relic plastid. However, identification of non-nuclear DNAs (*plDNA* and *mtDNA*) in the available *P. marinus* genome have proven elusive, perhaps due to the yet incomplete genome assembly and annotation. This process has been extremely challenging due to various factors, including the repetitiveness of the genome sequences and the uncertainty of this species' ploidy. In this regard, others and we have shown genetic variation in *Perkinsus* trophozoites that suggests their diploid status (Robledo et al., 1999; Reece et al., 2001; Thompson et al., 2011). The initial assembly of the *P. marinus* genome is considerably fragmented, composed by thousand of scaffolds (more than 17,000, some composed of a single contig) and many apparent genome duplications containing up to 90 genes. This is indicative of unresolved assemblies due to either multiple haplotypes or artifacts in the automated scaffolding. Although the genome annotation also contains a large proportion of “broken” gene structures lacking 5' or 3' ends, mining the genome with nuclear encoded apicoplast resident genes from *Plasmodium* as queries showed hits to numerous genes (Supplementary Table S1). The fact that we have not yet been able to identify a plastid genome or even a single gene encoding a plastid-targeted protein that is related to gene expression (e.g. ribosomal proteins) seems to suggest that *Perkinsus* lacks a plastid genome. This is also consistent with what we know about dinoflagellate plastid genomes, where virtually all genes not related to photosystems have been moved to the nucleus: it is easy to see how such an organelle could lose its genome once photosynthesis was lost. Other possibilities have yet to be formally ruled out, including the plastid DNA having been lost during the isolation of DNA (by spooling, to avoid shearing (Green, 1997)) for sequencing the genome (J.A. Fernández-Robledo, G.R. Vasta, unpublished results). The plastid DNA might also be represented by a low copy number and missed by

the current sequencing coverage. Deeper sequencing is needed to resolve this possibility. Although *Perkinsus* possesses a mitochondrial genome (Masuda et al., 2010), it is also absent in the current *Perkinsus* genome draft, opening the possibility that the plastid genome might have been excluded by the same bias.

3.2. Multimembraned structures in *Perkinsus*

Since its initial identification (Mackin et al., 1950), seven *Perkinsus* spp. have been described to date based on their geographic location, host species, morphology by light and transmission electron microscopy (TEM) and molecular analysis (Lester and Davis, 1981; Azevedo, 1989; Blackburn et al., 1998; McLaughlin et al., 2000; Coss et al., 2001a; Murrell et al., 2002; Casas et al., 2004; Burrenson et al., 2005; Dungan and Reece, 2006; Moss et al., 2008). However, even in those *Perkinsus* spp. described after the existence of the apicoplast was reported (Kilejian, 1974; Gardner et al., 1991; Howe, 1992; Williamson et al., 1994; McFadden et al., 1996; Wilson et al., 1996; Köhler et al., 1997), no plastid-like multimembrane organelles were noticed (McLaughlin et al., 2000; Coss et al., 2001a, b; Casas et al., 2004; Dungan and Reece, 2006; Moss et al., 2008). Recently, a plastid-like structure with four clearly defined membranes was observed in a single *Perkinsus atlanticus* (hereafter *Perkinsus olseni* (Murrell et al., 2002)) zoospore (Teles-Grilo et al., 2007b). However, the rigorous attribution of this structure to a bona fide plastid is doubtful because: (i) the putative plastid shows membranes that are further apart and appear to enclose an empty compartment; in general plastid membranes appear to be bound together by protein mass (Vaishnav and Striepen, 2006); (ii) the putative plastid is large (375–800 nm) and it is perplexing that it was not observed in the intact trophozoite; and (iii) the observed structure could be alternatively interpreted as a product of autophagy, a process that recycles macromolecules resulting from cellular remodelling or aberrant organelles (Reggiori and Klionsky, 2005; Yorimitsu and Klionsky, 2005). Similar structures with up to three membranes can be

identified in both trophozoites and zoospores from previously published work (e.g. Fig. 17 (Perkins, 1976), Fig. 19 (Azevedo, 1989), Fig. 20 (Coss et al., 2001b), Fig. 12 (Sunila et al., 2001)). Nevertheless, in our micrographs of *Perkinsus chesapeaki* (syn. *Perkinsus andrewsi*) zoospores, unique multimembrane structures that resemble the apicoplast in size can also be observed (Fig. 3). Although suggestive of the presence of a secondary plastid in *P. olseni* and *P. chesapeaki*, the morphological evidence is clearly insufficient to rigorously establish the presence of this organelle. To demonstrate this, the subcellular location of putative plastid-targeted proteins will need to be analysed by in situ immunofluorescence or immuno-electron microscopy, as was used to identify the plastid in apicomplexans (Waller et al., 1998).

3.3. Protein targeting to plastids

When the genes for plastid-derived proteins moved to the nucleus of the host during integration of a secondary plastid, they acquired targeting sequences to ensure their protein products are trafficked to the correct location within the cell. These sequences generally consist of a bipartite N-terminal extension composed of a canonical signal peptide (SP) followed by a transit peptide (TP) (Foth et al., 2003; Tonkin et al., 2008a, b). This mechanism is conserved to the point that has enabled the development of effective bioinformatic tools for prediction of apicoplast-targeted proteins based solely on primary sequence (e.g. a programme developed for predicting apicoplast resident genes in *Plasmodium* (<http://gecco.org.chemie.uni-frankfurt.de/pats/pats-index.php>) (Zuegge et al., 2001)). In addition, net charge and chaperone-binding sites are essential for robust targeting (Foth et al., 2003). It is noteworthy, however, that some proteins located in (or adjacent to) the outer apicoplast membrane lack any obvious common targeting motif (non-leader proteins) (Cavalier-Smith, 2003; Lim et al., 2009).

3.3.1. Does *P. marinus* superoxide dismutase 2 (PmSOD2) localise to multiple single membrane compartments?

Our previous work on PmSOD1 identified the presence of a leader sequence, suggesting mitochondrial targeting (Wright et al., 2002). IFA demonstrated that the protein co-localises with MitoTracker Red, a mitochondrial dye (Schott and Vasta, 2003). Hydrophathy analysis of the deduced amino acid sequence of a second SOD (PmSOD2) identified a strong hydrophobic region of approximately




Fig. 3. Ultrastructure of *Perkinsus chesapeaki* zoospores. A putative plastid has recently been identified in both zoospores and trophozoites of *Perkinsus olseni* (Teles-Grilo et al., 2007b). We have also identified putative plastid-like organelles in *P. chesapeaki* zoospores: A, plastid-like structure close to the nucleus (arrow); B and C, Detail of putative plastid-like organelles showing tightly associated membranes. N, nucleus; V, vacuole; F, flagellum; A, scale bar = 1 μ m; B, scale bar = 500 nm; C scale bar = 300 nm.




Fig. 4. Immunoelectron microscopy of *Perkinsus marinus* superoxide dismutase 2 (PmSOD2) in *P. marinus* trophozoites. Anti-recombinant PmSOD2 (rPmSOD2)-specific antibodies localised the gold grains to compartments (vacuoles) limited by single membranes (arrow). The thickness of these membranes (14 nm) was within the range of most eukaryotic membrane lipid bilayers. The gold particles appeared to localise to vesicle-like structures on the outer face of the membrane surrounding the vacuole, and either free or associated to amorphous material in the vacuole lumen, probably derived from the content of the above mentioned vesicles. Scale bar = 500 nm.

25 amino acid residues which was interpreted as a membrane-spanning domain of type II transmembrane proteins (Wright et al., 2002) and later proposed as suggestive of the presence of a relict plastid in *Perkinsus* (Saldarriaga et al., 2004). IFA revealed, however, that PmSOD2 localises to multiple subcellular structures/organelles (Schott and Vasta, 2003). Immunogold electron microscopy (IEM) using a polyclonal antibody against recombinant PmSOD2 outlines multiple vesicles containing clustered gold grains and electron-dense material of unknown nature, both of which appear to be released into a large single membrane compartment (Fernández-Robledo et al., 2008b) (Fig. 4). Transfection of *Perkinsus* trophozoites with pPmSOD2(MOE)-GFP resulted in PmSOD2-GFP localised to multiple small compartments and to what appeared as a single large compartment, supporting the hypothesis that the small vesicles visualised by the gold grains might either fuse and/or empty their contents in a larger compartment (Fernández-Robledo et al., 2008b). However, we cannot rule out that the protein was either in transit as endoplasmic reticulum (ER) vesicles to its final destination, the trophozoite was dividing or simply mislocalised due to the use of the non-native PmSOD2 flanking regions. We have observed that disruption of the flanking regions in the transfection vector may result in gene deregulation (J.A. Fernández-Robledo, G.R. Vasta, unpublished results). In either case, the localisation data are not consistent with PmSOD2 being a plastid protein.

3.3.2. *ispC* (1-deoxy-d-xylulose 5-phosphate reductoisomerase) also localises to multiple compartments

Some enzymes of the MEP pathway for the synthesis of isoprenoids have been experimentally confirmed as associated with the apicoplast of apicomplexans (Jomaa et al., 1999; Ralph et al., 2004b). All six MEP enzymes and *ispD*, which was missed in previous genome mining and found in *Perkinsus*, have bipartite targeting peptides at the N-terminus (Matsuzaki et al., 2008) which is characteristic of proteins targeted to secondary plastids, thereby supporting the idea that *Perkinsus* has a non-photosynthetic secondary plastid. More intriguingly, immunofluorescence microscopy of the 1-deoxy-D-xylulose 5-phosphate reductoisomerase (*IspC*) reveals multiple small compartments suggesting that *Perkinsus* might have multiple plastids, even though the compartments do not appear to contain detectable amounts of DNA (Matsuzaki et al., 2008). GFP labelling of *IspC* under MOE gene control also highlights several structures per trophozoite (M. Matsuzaki, H. Kuroiwa, H. Nozaki, T. Nozaki and K. Kita, unpublished data).

3.4. Presence in *P. marinus* of other pathways/structures typically associated with plastids

3.4.1. Type II FAS enzymes

Driven by the lack of a chemotherapeutic treatment to reduce or eliminate *P. marinus* infections in oysters and their interest in lipid metabolism (Chu et al., 2002, 2004; Lund and Chu, 2002; Lund et al., 2007), Chu's group identified Triclosan, a specific inhibitor of FabI, which is a Type II FAS enzyme found in prokaryotes and plastids, as an inhibitor of *P. marinus* proliferation (Lund et al., 2005). The Triclosan results, however, should be interpreted with caution as revealing strictly plastid-associated FASII biosynthesis, since the apicomplexan *Theileria* lacks FASII genes (Gardner et al., 2005) and yet is susceptible to this drug. Furthermore, in *Plasmodium*, FabI is not the target of the antimalarial activity of Triclosan (Yu et al., 2008). Nevertheless, *Perkinsus* is sensitive to inhibitors of other typically plastid-localised enzymes involved in fatty acid biosynthesis (e.g. acetyl-CoA carboxylase) (Stelter et al., 2007). Unlike most eukaryotes, the FAS II-dependent lipoic acid synthesis has been associated with the apicoplast in both *Plasmodium falciparum* and *Toxoplasma gondii* (reviewed in Seeber and Soldati-Favre,

2010), and we have identified a putative lipoate synthase (XP_002786331) in *Perkinsus*, although predictions for its localisation are inconclusive.

3.4.2. Fe-S cluster biosynthesis

Fe-S clusters are ubiquitous prosthetic groups required to sustain fundamental life processes including electron transfer, substrate binding/activation, Fe-S storage, regulation of gene expression and regulation of enzyme activity (Xu and Moller, 2011). In addition to the mitochondrial Fe-S cluster biosynthesis, plastid-harboured eukaryotes usually have a second assembly for the Fe-S cluster biosynthesis in the plastid (Fleige et al., 2010; Kalanon and McFadden, 2010; Lim and McFadden, 2010), including the apicoplast (Seeber and Soldati-Favre, 2010). The identification of transcripts of the plastid-type ferredoxin and its associated reductase in in vitro-cultivated *P. marinus* has also been proposed as indicative of the presence of a plastid since this redox pair is exclusively found in cyanobacteria and plastid-harboured organisms (Stelter et al., 2007). Other Fe-S cluster genes predicted to localise to the relic plastid include PmSufB (see Section 4.2).

3.4.3. Haeme synthesis

In most eukaryotes, haeme is an important prosthetic group on many proteins, such as cytochromes and peroxidases, which are involved in electron transfer and redox chemistry. In photosynthetic organisms (plants/algae), the complete pathway for tetrapyrrole biosynthesis is harboured within the plastid whereas in animals/fungi the pathway is distributed between the cytosol and mitochondria. In *Plasmodium*, a hybrid pathway runs cooperatively involving the cytosol, mitochondria and the apicoplast, that not only fulfil the parasite's requirements for the above processes, but also those related to the heme derived from the haemoglobin in the host erythrocytes (reviewed in Lim and McFadden, 2010; Nagaraj et al., 2010). The PBGS (as represented by *T. gondii* and *P. falciparum* counterparts) is conserved across the phyla and is very similar to the plant enzyme in its biochemical properties (Jaffe et al., 2011). Key features in the primary sequence of the protein help to identify it as the plant type rather than the animal type of enzyme. A comparison of the PBGS enzymes from *P. marinus* and *T. gondii* and several other PBGS reveals that, unlike the apicomplexan enzyme, the *Perkinsus* enzyme (XP_002782704) is distinctly of the animal type and probably located in the cytosol rather than the plastid. There is precedence for the co-existence of animal and plant type pathways for heme biosynthesis in a single cell (e.g. *Euglena gracilis*) and it is probable that an apicomplexan ancestor contained an animal type pathway to begin with and acquired the plant type pathway via the secondary endosymbiotic event. A brief period must have existed where the apicomplexan ancestor contained both pathways, after which the different lineages developed a mosaic of both pathways. According to this hypothesis, if the common ancestor for the Apicomplexa, Chromerida, Dinophyta and Perkinsozoa contained both the animal and plant pathways, it appears that (as of now based on PBGS only), all except Perkinsozoa have retained the plant type pathway (as a consequence of the presence of the plastid in all of them). This observation alone is not sufficient to suggest that *Perkinsus* might lack a plastid as some apicomplexans have totally lost the pathway for heme biosynthesis (e.g. *Cryptosporidium* (in this case the plastid itself is lost) and *Theileria*) (Abrahamsen et al., 2004; Gardner et al., 2005; Pain et al., 2005). Therefore it is also possible that *Perkinsus* has the plastid but harbours a complete animal type pathway for heme biosynthesis. Further characterisation is needed to characterise the plastid-like heme biosynthesis pathway.

3.4.4. ABA

ABA is a stress hormone common in plants (Nambara and Marion-Poll, 2005). More recently, it has been identified in *Toxoplasma*, where it is involved in Ca^{2+} -mediated egress from the host cell (Nagamune et al., 2008a, b). *Toxoplasma* is sensitive to fluridone, an herbicide that inhibits the indirect pathway for ABA synthesis; treatment of *Toxoplasma* cultures with fluridone resulted in both delayed egress and induction of cysts (Nagamune et al., 2008a). In plants, ABA synthesis uses an indirect pathway that involves β -carotene and resides in the plastid (Schwartz et al., 2003). In *P. falciparum* a dual localisation in the mitochondria and apicoplast has been proposed but details remain unresolved (Tonhosolo et al., 2009). Indeed, *Perkinsus* is sensitive to the herbicide fluridone (J.A. Fernández-Robledo, G.R. Vasta, unpublished results), which targets the enzyme phytoene dehydrogenase (XP_002772671); in addition, *Perkinsus* has at least three classes of genes carrying the G protein-coupled receptor 89A domain (XP_002783768, XP_002775994, XP_002775137).

3.4.5. Shikimate pathway

Shikimate and chorismate are precursors in the biosynthesis of aromatic amino acids and other aromatic secondary metabolites in microorganisms and plants. Recently, the presence of a shikimate pathway in *P. olsenii* was demonstrated by inhibition studies with glyphosate, a potent inhibitor of 5-enolpyruvylshikimate-3-phosphate synthase, a key enzyme in the shikimate pathway. Glyphosate inhibited the in vitro proliferation of *P. olsenii* in a dose-dependent manner, and this effect was reversed by addition of chorismate (Elandalloussi et al., 2005, 2008). Although in Apicomplexa this pathway is cytosolic, in dinoflagellates it localises to the plastid (Waller et al., 2006). In the *Perkinsus* genome, we have identified several genes of the shikimate pathway, including chorismate synthase (XP_002773541) and a gene (XP_002773648) with a match to the pentafunctional arom protein, which catalyses five consecutive enzymatic reactions in prechorismate polyaromatic amino acid biosynthesis,

3.4.6. ER-associated protein degradation (ERAD) system

The ERAD pathway is aimed at tagging misfolded proteins of the ER for ubiquitination and subsequent degradation, and participates in the import of apicoplast proteins (Sheiner and Soldati-Favre, 2008; Lim et al., 2009; Spork et al., 2009). Searching for the core components of the ERAD in the *Perkinsus* genome, we have identified Cdc48, Ufd1 and Npl4. The *Perkinsus* genome also has numerous Derin homologues (intramembrane serine proteases that cleave substrates in or near transmembrane domains) necessary for processing proteins into organelles. Similar to apicomplexan parasites (Agrawal et al., 2009), the presence of multiple copies of some of these genes in *Perkinsus* opens the possibility that some of them might reside in the putative plastid, although we must note that these proteins are also expected to exist in the absence of a plastid and consequently the identification of specific copies related to plastid-targeting will require direct experimental evidence.

4. Working hypothesis and outstanding questions

Under the Chromalveolata hypothesis, some extant non-photosynthetic lineages, including *Perkinsus*, must have derived from photosynthetic ancestors in which the plastid membranes have evolved into diverse structures (Keeling, 2010). In this regard, membranes may arise by growth and division from pre-existing membranes (designated as “genetic membranes”) or by differentiation from another type of membrane (“non-genetic membranes”, e.g. lysosome membranes) (Cavalier-Smith, 1995, 2003). Thus, membranes




Fig. 5. Working hypotheses for the presence of a plastid in the protozoan *Perkinsus*. (A) A secondary endosymbiosis resulted in an ancestor plastid with multiple membranes and a plastid genome. (B) *Perkinsus* adaptation to intracellular parasitism resulted in plastid size reduction, photosystem gene loss, gene transfer to the nucleus (N) and acquisition of bipartite signal signatures for re-targeting proteins to the plastid. (C) Hypothetical targeting of *Perkinsus* nuclear-encoded proteins to the relic plastid via the endoplasmic reticulum (ER), based on the most parsimonious *Plasmodium* model (Tonkin et al., 2008b). MEP, non-mevalonate; FAS, fatty acid synthesis; Fe-S, iron-Sulphur; ABA, abscisic acid; SP, signal peptide; TP, transit peptide; MP, mature peptide.

budded from either the host Golgi or the ER might have been added to those from the original endosymbiont (Cavalier-Smith, 2003). As a result, the protein-targeting and translocation machinery has been modelled by the natural history of any particular secondary endosymbiosis within a lineage and lead to increased complexity in the host cell. Based on the fragmentary evidence described above, two working hypotheses can be posed for the presence of a relic plastid in *Perkinsus* (Fig. 5). In hypothesis 1 *Perkinsus* shares with other Alveolata a relic plastid with multiple membranes and a plastid genome. During cell propagation the organelle is segregated, unless sexual stages are confirmed, in which case it may occur that only one gamete inherits it. In hypothesis 2 *Perkinsus* has lost the plastid genome but the plastid organelle has been retained and is nevertheless segregated in zoospores and trophozoites. These hypotheses are discussed in the following sections.

4.1. Is there physical evidence for a multimembrane structure in *Perkinsus* trophozoites or zoospores?

Serial TEM images and three-dimensional (3D) reconstruction have proved useful to measure organellar volumes, geometries and subcellular features, and have enabled the development of 3D cellular models in protozoan parasites of sizes comparable to *Perkinsus* life stages (Hopkins et al., 1999; Elliott, 2007; Elliott et al., 2008). The application of similar serial TEM images and 3D reconstruction to *Perkinsus* spp. would provide a precise blueprint of its subcellular organisation and based on the reported *Perkinsus* cell size and that of the reported relic plastid (Teles-Grilo et al., 2007b), 5–10 sections should reveal the putative organelle (Elliott, 2007). The cryo tomographic X-ray techniques that, in protozoans, have enabled the detailed survey of nucleus, digestive vacuoles and membrane networks in an intact cellular context (Hanssen et al., 2010, 2011), have also been useful to show integration of the apicoplast with other organelles (Kudryashev et al., 2010; Miranda et al., 2010). Application of this approach to *P. marinus* represents a promising avenue for the high resolution of yet unidentified subcellular structures. Further, a putative plastid genome has not yet been identified in any member of the Perkinsozoa and DNA dyes fail to stain any subcellular structure in the trophozoite other than the nucleus and mitochondrion (Matsuzaki et al., 2008). There is no indication from the genome project for a plastid genome but neither is there data from the mitochondrion despite the fact that a partial *Perkinsus* mitochondrial genome has been reported by other means (Masuda et al., 2010). Thus, in addition to further mining of the genome, alternative approaches for identifying the plastid genome should include DNA staining by 5-bromodeoxyuridine (BrdU) incorporation followed by IEM or fluorescently-labelled anti-BrdU antibodies. This technique combined with IEM staining of alternate sections for genes predicted as targeted to the relic plastid (e.g. PmSOD2 or IspC) should reveal whether any other compartments contain DNA and potentially represent the relic plastid. Putative tRNAs, rRNAs and ORFs typically found on plastid genomes can also be used in high resolution in situ hybridisation to localise the putative plastidic transcripts (McFadden et al., 1996) or in pulse field gel electrophoresis of *Perkinsus* cells (Teles-Grilo et al., 2007a) optimised for the size range of *pl*DNA. Usefulness of the latter approach might be hard to predict if, similar to dinoflagellates, the *Perkinsus* plastid genome is in 2–3 Kb circles (Williams and Keeling, 2003; van der Giezen et al., 2005; Howe et al., 2008; Slamovits and Keeling, 2008; Hjort et al., 2010; Keeling, 2010).

4.2. What is the subcellular destination for the *Perkinsus* proteins containing bipartite signal sequences?

If no plastid genome exists, the organelle can still be identified by localising nucleus-encoded plastid-targeted proteins. To date, 10 genes have been identified in *Perkinsus* that either have a bipartite signal or encode products that are expected to target the putative plastid. These include ferredoxin (*PmFD*) and its associated reductase (*PmFNR*), acetyl-CoA carboxylase and the six genes from the MEP pathway (*dxs*, *ispC*, *ispE*, *ispF*, *ispG*, *ispH*) (Matsuzaki et al., 2006, 2008; Grauvogel et al., 2007; Stelter et al., 2007; Joseph et al., 2010). In addition, the use of Prediction of Apicoplast Targeting Sequences (PATS) (Zuegge et al., 2001) resulted in numerous putative *Perkinsus* plastid genes, including most of the genes already identified in the literature (Supplementary Table S2), some of them carrying the predicted bipartite signal (e.g. PmSufB, Fig. 6). The bipartite extension of apicomplexan proteins is well established as diagnostic for targeting to secondary plastids (Foth et al., 2003; Ralph et al., 2004a; Tonkin et al., 2006a, b, 2008a, b), however the rigorous assignment of a specific subcellular localisation

for any *Perkinsus* putative plastid gene will require experimental validation.

5. Technical considerations

5.1. Species and strains

Although six out of the seven *Perkinsus* spp. are available at public repositories such as the American Type Culture Collection (ATCC), USA, <http://www.atcc.org/>), structural and genetic evidence for the presence of a plastid have only been obtained in three species/strains. These include *P. marinus* CB5D4 (ATCC PRA-240), the strain used for the sequence of the genome and transcriptome (Fernández-Robledo et al., 2008a; Joseph et al., 2010), *P. olseni* ALG1 (Robledo et al., 2002), the species where the relic plastid was described (Teles-Grilo et al., 2007b), and *P. chesapeakei* A8–4a (ATCC #50807) (= *P. andrewsi*), which spontaneously sporulate in culture (Coss et al., 2001a, b; Burrenson et al., 2005) and for which we have identified plastid-like structures in the zoospores (Fig. 3).

5.2. Comparative genomics and proteomics

The parallel implementation of state-of-the art deep DNA sequencing methodology to *Perkinsus* spp. such as *P. marinus* and *P. chesapeakei* will contribute enormously to a more rigorous search, annotation and curation of putative plastid genes and the search for the plastid genome (possibly with a codon bias or signature) in *Perkinsus* by yielding less partitioned and redundant assemblages that also include clearly identified data from the mitochondrial genome. Further, the identification of *Perkinsus* homologues of proteins targeted to the apicoplast should provide bases for identifying the organelle where proteins with bipartite-N-terminal signal ultimately function. Whole-genome analysis of unicellular eukaryotes can also reveal those gene loss and gain events often associated with the lifestyle adopted by each particular group of organisms (Martens et al., 2008). Furthermore, deep sequencing should provide enough resolution to determine whether *P. marinus* and *P. chesapeakei* are distinct bona fide *Perkinsus* spp. or genetically distinct *P. marinus* assemblages. The direct comparison of putative plastid genes in the *P. marinus* and *P. chesapeakei* genomes might also shed light on the genetic/molecular basis of why the latter spontaneously sporulates in culture whilst the former does not. Similar to apicomplexans (Lim et al., 2009), a well-curated *Perkinsus* genome will further enable the application of proteomic approaches for the identification of putative genes that may reside in the relic plastid whilst lacking the canonical bipartite signal sequence.

5.3. Cellular and molecular tools, and surrogate models to study *Perkinsus* plastid targeting

Success in the search for the *Perkinsus* plastid will require availability of organelle and structural markers. Few commercial markers have been used in *Perkinsus* other than Mitotracker for mitochondria and nuclear stains (DAPI, Sybr Green). However, multiple fluorescent dyes are available (Molecular Probes, Invitrogen) that could be tested and validated in *Perkinsus*. Furthermore, our *Perkinsus* genome and transcriptome projects have enabled the identification of numerous genes predicted as targeted to most organelles and subcellular structures (Table 1, Supplementary Table S1) which, combined with the availability of numerous spectral variants of fluorescent proteins and epitope tags, should enable multicolour imaging and co-localisation studies using confocal microscopy. The development of subcellular markers for *Perkinsus* should also represent a useful tool for the scientific community




Fig. 6. Clustal W alignment of *Perkinsus marinus* putative iron-Sulphur (Fe-S) assembly ATPase SufB. The box above the alignment displays the results of Sliding Window Iteration of TargetP (SWIT) (Matsuzaki et al., 2008) that shows the distribution of targeting preferences for the N-terminal pre-sequences. The neural network score (y axis) of the TargetP prediction is plotted for the N-terminal 150 amino acids (x axis). Although the signal peptide (SP) prediction is not very strong, the chloroplast transit peptide (cTP) is predicted to be near the SP cleavage site and, therefore, as a good candidate for a class II bipartite leader. Green, plastid target peptide; light blue: mitochondrial target peptide; yellow, signal peptide; grey, other locations. *Thiomicrospira*: *Thiomicrospira crunogena*; *Vibrio*: *Vibrio caribbenthicus*; *Roseobacter*: *Roseobacter litoralis*; *Ketogulonicigenium*: *Ketogulonicigenium vulgare*; *Perkinsus*: *P. marinus*.

interested in the Chromalveolata. The sizes of the *P. marinus* trophozoites (2.7–10 μm; 30–80 μm prior to zoosporulation) and zoospores (2–3 × 4–6 μm) are similar to those equivalent life stages from the apicomplexan parasites for which this technique has demonstrated sufficient resolution to address questions about secretion and cell division through the use of subcellular markers (Vaishnav and Striepen, 2006; Nishi et al., 2008; Sheiner and Soldati-Favre, 2008). There is mounting evidence to suggest that *Perkinsus* uses transplicing (Zhang et al., 2007; Joseph et al., 2010; Lin et al., 2011), a process where mRNA can carry a splice-leader, which is an indication of post-transcriptional regulation (Zhang et al., 2011). Since it remains unknown whether RNA interference (RNAi) (EER05359.1, EER19214.1) plays a role in these processes, coding sequence of the genes of interest should be cloned under the native flanking regions to avoid over-expression artifacts. Finally, since studies on plastid targeting in apicomplexans (Foth et al., 2003; Ralph et al., 2004a; Tonkin et al., 2006a, b, 2008a, b) have revealed that these mechanisms are very well conserved, swapping bona fide apicoplast genes with the putative *Perkinsus* homologues can be used to test whether the *Perkinsus* relic plastid and the apicoplast were similarly shaped by a similar intracellular parasitic lifestyle. For example, *T. gondii*, a well-developed genetic model with multiple tools available (Ajioka and Soldati, 2007; Weiss and Kim, 2007) may be suitable as a surrogate model to examine function and subcellular targeting of *Perkinsus* proteins.

6. Concluding remarks

The several lines of evidence described above (homologues of type II FAS, Fe-S cluster, heme, ABA, lipoic acid, shikimate and other pathway components) suggest the presence of plastid-targeted genes in *Perkinsus* spp. Although the enzymes involved in these pathways are clearly predicted to target the putative plastid, further experimental characterisation and localisation will be required. In this regard, intriguing subcellular structures noticed in earlier studies raise interesting questions about whether *Perkinsus* has a relic plastid and, if so, the number of membranes that surround it. As the first described and better characterised member of the Perkinsozoa, research on *Perkinsus* plastid localisation, targeting and biogenesis will provide fundamental insights into the biology of this phylum. Further, given its phylogenetic position within the Alveolata, the characterisation of the putative plastid genes and the potential relic organelle to which the products are targeted, should make this parasite a promising model organism for investigating various aspects of plastid biology (e.g. biogenesis, segregation, targeting, translocation, functional regulation) in an evolutionary context. Analysis of the targeting peptides will lead to the identification and functional characterisation of the plastid-targeted proteins, which may reveal *Perkinsus* adaptations to intracellular survival, proliferation and virulence. If the absence of a multimembrane compartment is established, *Perkinsus* will

Table 1Proposed *Perkinsus* genes for co-localisation studies derived from published studies or from genome mining.

Organelle	Marker	Accession	Reference
Nucleus	Histone H3	XM_002788843	<i>Perkinsus</i> genome
	Dmc1	XM_002784146	
Centrosome	Centrin	XM_002787551	<i>Perkinsus</i> genome
Golgi	Ras	XM_002786157	<i>Perkinsus</i> genome
	Rab6	XM_002788779	
	Sys1	XM_002781689	
ERAD	Cdc48	XM_002782252	<i>Perkinsus</i> genome
	Ufd1	XM_002786555	
	Npl4	XM_002778993	
Membranes/vesicles	SNARE	XP_002784625	<i>Perkinsus</i> genome
	SNAP	XP_002769618	
ER	KDEL	Pmar PMAR017428	<i>Perkinsus</i> genome
Mitochondria	PmSOD1	AY095212	Schott and Vasta (2003)
	Cit c oxydase	AB513789	Masuda et al. (2010)
Putative plastid	MEP pathway		Matsuzaki et al. (2008)
	Fe-S cluster pathway		Stelter et al. (2007)
Large vacuole	PmAPX2	XP_002767285	Schott et al. (2004)
Cell wall	PmMOE?	EF632302	Fernández-Robledo et al. (2008a)
	GH18 Chitinase	XP_002768327	<i>Perkinsus</i> genome
Cytoskeleton	Actin	AY436364	Reece et al. (1997)
	Tubulin	XM_002788606	<i>Perkinsus</i> genome
Cytosol	Glycolytic enzymes	XM_002788328	<i>Perkinsus</i> genome
		XM_002783157	Joseph et al. (2010)

be useful to gain insight into molecular innovations that compensate for plastid loss in intracellular parasites, an event that under the Chromalveolata hypothesis has taken place at multiple points during evolution. *Perkinsus marinus* is virtually ubiquitous along the Atlantic and Gulf coasts of the USA, and has devastated natural and farmed oyster populations. No effective therapeutic intervention strategies against *Perkinsus* infections are available at the present time and the impact of the disease in the environment is hardly contained by the current management practices. Soon after the hatchery-raised seed is deployed in the environment, it becomes infected and the oyster farmers harvest their product as soon as it reaches commercial size, before *Perkinsus* infections cause death. Demonstrating the presence of a relic plastid in *Perkinsus* and characterising its biology may lead to potential targets for the development of chemotherapeutic drugs for use in contained oyster farming settings or oysters genetically modified for disease resistance. Furthermore, a deep understanding of the biology of the putative plastid genes in *Perkinsus* will complement ongoing studies on apicomplast genes, thereby contributing to a better understanding of cellular mechanisms that might result in the development of novel therapeutic targets against apicomplexan parasites.

Acknowledgements

This study was supported by Grant 1R21AI076797-01A2 from the National Institutes of Health (NIH), USA, partially by Grant 0333240 from National Science Foundation (NSF)/United States Department of Agriculture (USDA)-CSREES NSF/USDA, and by Grants IOB 0618409, IOS 0822257 and IOS 0958016 from NSF.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.ijpara.2011.07.008.

References

- Abrahamsen, M.S., Templeton, T.J., Enomoto, S., Abrahante, J.E., Zhu, G., Lancto, C.A., Deng, M., Liu, C., Widmer, G., Tzipori, S., Buck, G.A., Xu, P., Bankier, A.T., Dear, P.H., Konfortov, B.A., Spriggs, H.F., Iyer, L., Anantharaman, V., Aravind, L., Kapur, V., 2004. Complete genome sequence of the apicomplexan, *Cryptosporidium parvum*. *Science* 304, 441–445.
- Agrawal, S., van Dooren, G.G., Beatty, W.L., Striepen, B., 2009. Genetic evidence that an endosymbiont-derived endoplasmic reticulum-associated protein degradation (ERAD) system functions in import of apicomplast proteins. *J. Biol. Chem.* 284, 33683–33691.
- Ajioka, J.W., Soldati, D., 2007. *Toxoplasma*: Molecular and Cellular Biology. Horizon Bioscience, Norfolk, UK.
- Azevedo, C., 1989. Fine structure of *Perkinsus atlanticus* n. sp. (Apicomplexa, Perkinsea) parasite of the clam *Ruditapes decussatus* from Portugal. *J. Parasitol.* 75, 627–635.
- Bachvaroff, T.R., Handy, S.M., Place, A.R., Delwiche, C.F., 2011. Alveolate phylogeny inferred using concatenated ribosomal proteins. *J. Eukaryot. Microbiol.* 58, 223–233.
- Bhattacharya, D., Yoon, H.S., Hackett, J.D., 2004. Photosynthetic eukaryotes unite: endosymbiosis connects the dots. *BioEssays* 26, 50–60.
- Blackbourn, J., Bower, S.M., Meyer, G.R., 1998. *Perkinsus qugwadi* sp.nov. (*incertae sedis*), a pathogenic protozoan parasite of Japanese scallops, *Patinopecten yessoensis*, cultured in British Columbia, Canada. *Can. J. Zool. Rev. Can. Zool.* 76, 942–953.
- Brugerolle, G., 2002. *Cryptophagus subtilis*: a new parasite of cryptophytes affiliated with the Perkinsozoa lineage. *Eur. J. Protistol.* 37, 379–390.
- Burreson, E.M., Reece, K.S., Dungan, C.F., 2005. Molecular, morphological, and experimental evidence support the synonymy of *Perkinsus chesapeakei* and *Perkinsus andrewsi*. *J. Eukaryot. Microbiol.* 52, 258–270.
- Bushek, D., Ford, S.E., Chintala, M.M., 2002. Comparison of *in vitro*-cultured and wild-type *Perkinsus marinus*. III. Fecal elimination and its role in transmission. *Dis. Aquat. Organ.* 51, 217–225.
- Casas, S.M., Grau, A., Reece, K.S., Apakupakul, K., Azevedo, C., Villalba, A., 2004. *Perkinsus mediterraneus* n. sp., a protistan parasite of the European flat oyster *Ostrea edulis* from the Balearic Islands, Mediterranean Sea. *Dis. Aquat. Organ.* 58, 231–244.
- Cavalier-Smith, T., 1995. Membrane heredity, symbiogenesis, and the multiple origins of algae. In: Arai, R., Kato, M., Doi, Y. (Eds.), *Biodiversity and Evolution*. The National Science Museum Foundation, Tokyo, pp. 75–114.
- Cavalier-Smith, T., 1998. A revised six-kingdom system of life. *Biol. Rev. Camb. Philos. Soc.* 73, 203–266.
- Cavalier-Smith, T., 1999. Principles of protein and lipid targeting in secondary symbiogenesis: euglenoid, dinoflagellate, and sporozoan plastid origins and the eukaryote family tree. *J. Eukaryot. Microbiol.* 46, 347–366.
- Cavalier-Smith, T., 2003. Genomic reduction and evolution of novel genetic membranes and protein-targeting machinery in eukaryote-eukaryote chimaeras (meta-algae). *Philos. Trans. R. Soc. Lond. B.* 358, 109–134.

- Chu, F.L., Lund, E., Soudant, P., Harvey, E., 2002. *De novo* arachidonic acid synthesis in *Perkinsus marinus*, a protozoan parasite of the eastern oyster *Crassostrea virginica*. *Mol. Biochem. Parasitol.* 119, 179–190.
- Chu, F.L., Lund, E.D., Harvey, E., Adlof, R., 2004. Arachidonic acid synthetic pathways of the oyster protozoan parasite, *Perkinsus marinus*: evidence for usage of a delta-8 pathway. *Mol. Biochem. Parasitol.* 133, 45–51.
- Coss, C.A., Robledo, J.A., Ruiz, G.M., Vasta, G.R., 2001a. Description of *Perkinsus andrewsi* n. sp. isolated from the Baltic clam (*Macoma balthica*) by characterization of the ribosomal RNA locus, and development of a species-specific PCR-based diagnostic assay. *J. Eukaryot. Microbiol.* 48, 52–61.
- Coss, C.A., Robledo, J.A., Vasta, G.R., 2001b. Fine structure of clonally propagated *in vitro* life stages of a *Perkinsus* sp. isolated from the Baltic clam *Macoma balthica*. *J. Eukaryot. Microbiol.* 48, 38–51.
- Dungan, C.F., Reece, K.S., 2006. *In vitro* propagation of two *Perkinsus* spp. parasites from Japanese Manila clams *Venerupis philippinarum* and description of *Perkinsus honshuensis* n. sp. *J. Eukaryot. Microbiol.* 53, 316–326.
- Elandaloussi, L.M., Leite, R.B., Rodrigues, P.M., Afonso, R., Cancela, M.L., 2008. Effect of the herbicide Roundup on *Perkinsus olseni* *in vitro* proliferation and *in vivo* survival when infecting a permissive host, the clam *Ruditapes decussatus*. *Bull. Environ. Contam. Toxicol.* 80, 512–515.
- Elandaloussi, L.M., Rodrigues, P.M., Afonso, R., Leite, R.B., Nunes, P.A., Cancela, M.L., 2005. Shikimate and folate pathways in the protozoan parasite, *Perkinsus olseni*. *Mol. Biochem. Parasitol.* 142, 106–109.
- Elliott, D.A., 2007. Serial Sectioning via Microtomy. *Microscopy Today* 15, 30–33.
- Elliott, D.A., McIntosh, M.T., Hosgood 3rd, H.D., Chen, S., Zhang, G., Baevova, P., Joiner, K.A., 2008. Four distinct pathways of hemoglobin uptake in the malaria parasite *Plasmodium falciparum*. *Proc. Natl Acad. Sci. USA* 105, 2463–2468.
- Ellis, J.T., Morrison, D.A., Jeffries, A.C., 1998. The phylum Apicomplexa: an update on the molecular phylogeny. Kluwer Academic Publishers, Boston.
- Feagin, J.E., Parsons, M., 2007. The Apicoplast and Mitochondrion of *Toxoplasma gondii*. In: Weiss, L.M., Kim, K. (Eds.), *Toxoplasma gondii: the model apicomplexan: perspectives and methods*. Elsevier, San Diego, CA, USA, pp. 207–236.
- Fernández-Robledo, J.A., Lin, Z., Vasta, G.R., 2008a. Transfection of the protozoan parasite *Perkinsus marinus*. *Mol. Biochem. Parasitol.* 157, 44–53.
- Fernández-Robledo, J.A., Schott, E.J., Vasta, G.R., 2008b. *Perkinsus marinus* superoxide dismutase 2 (PmSOD2) localizes to single-membrane subcellular compartments. *Biochem. Biophys. Res. Commun.* 375, 215–219.
- Fichera, M.E., Roos, D.S., 1997. A plastid organelle as a drug target in apicomplexan parasites. *Nature* 390, 407–409.
- Figuerola, R.I., Garces, E., Massana, R., Camp, J., 2008. Description, host-specificity, and strain selectivity of the dinoflagellate parasite *Parvilicifera sinerae* sp. nov. (Perkinsozoa). *Protist* 159, 563–578.
- Fleige, T., Limentakis, J., Soldati-Favre, D., 2010. Apicoplast: keep it or leave it. *Microbes Infect.* 12, 253–262.
- Ford, S.E., 1996. Range extension by the oyster parasite *Perkinsus marinus* into the northeastern United States: Response to climate change? *J. Shellfish Res.* 15, 45–56.
- Ford, S.E., Chintala, M.M., 2006. Northward expansion of a marine parasite: Testing the role of temperature adaptation. *J. Exp. Mar. Biol. Ecol.* 339, 226–235.
- Ford, S.E., Smolowitz, R., 2007. Infection dynamics of an oyster parasite in its newly expanded range. *Mar. Biol.* 151, 119–133.
- Foth, B.J., Ralph, S.A., Tonkin, C.J., Struck, N.S., Fraunholz, M., Roos, D.S., Cowman, A.F., McFadden, G.I., 2003. Dissecting apicoplast targeting in the malaria parasite *Plasmodium falciparum*. *Science* 299, 705–708.
- Gardner, M.J., Bishop, R., Shah, T., de Villiers, E.P., Carlton, J.M., Hall, N., Ren, Q., Paulsen, I.T., Pain, A., Berriman, M., Wilson, R.J., Sato, S., Ralph, S.A., Mann, D.J., Xiong, Z., Shallom, S.J., Weidman, J., Jiang, L., Lynn, J., Weaver, B., Shoaibi, A., Domingo, A.R., Wasawo, D., Crabtree, J., Wortman, J.R., Haas, B., Angiuoli, S.V., Creasy, T.H., Lu, C., Suh, B., Silva, J.C., Utterback, T.R., Feldblyum, T.V., Perete, M., Allen, J., Nieman, W.C., Taracha, E.L., Salzberg, S.L., White, O.R., Fitzhugh, H.A., Morzaria, S., Venter, J.C., Fraser, C.M., Nene, V., 2005. Genome sequence of *Theileria parva*, a bovine pathogen that transforms lymphocytes. *Science* 309, 134–137.
- Gardner, M.J., Williamson, D.H., Wilson, R.J., 1991. A circular DNA in malaria parasites encodes an RNA polymerase like that of prokaryotes and chloroplasts. *Mol. Biochem. Parasitol.* 44, 115–123.
- Gauthier, J.D., Feig, B., Vasta, G.R., 1995. Effect of fetal bovine serum glycoproteins on the *in vitro* proliferation of the oyster parasite *Perkinsus marinus*: Development of a fully defined medium. *J. Eukaryot. Microbiol.* 42, 307–313.
- Gile, G.H., Patron, N.J., Keeling, P.J., 2006. EFL GTPase in cryptomonads and the distribution of EFL and EF-1alpha in chromalveolates. *Protist* 157, 435–444.
- Goggin, C.L., Barker, S.C., 1993. Phylogenetic position of the genus *Perkinsus* (Protista, Apicomplexa) based on small subunit ribosomal RNA. *Mol. Biochem. Parasitol.* 60, 65–70.
- Grauvogel, C., Reece, K.S., Brinkmann, H., Petersen, J., 2007. Plastid isoprenoid metabolism in the oyster parasite *Perkinsus marinus* connects dinoflagellates and malaria pathogens—new impetus for studying alveolates. *J. Mol. Evol.* 65, 725–729.
- Green, B.R., 2011. Chloroplast genomes of photosynthetic eukaryotes. *Plant J.* 66, 34–44.
- Green, E.D., 1997. *Genome analysis: a laboratory manual*. Cold Spring Harbor Laboratory Press, Plainview, New York.
- Hanssen, E., Carlton, P., Deed, S., Klonis, N., Sedat, J., DeRisi, J., Tilley, L., 2010. Whole cell imaging reveals novel modular features of the exomembrane system of the malaria parasite, *Plasmodium falciparum*. *Int. J. Parasitol.* 40, 123–134.
- Hanssen, E., Knoechel, C., Klonis, N., Abu-Bakar, N., Deed, S., LeGros, M., Larabell, C., Tilley, L., 2011. Cryo transmission X-ray imaging of the malaria parasite, *P. falciparum*. *J. Struct. Biol.* 173, 161–168.
- Hjort, K., Goldberg, A.V., Tsaousis, A.D., Hirt, R.P., Embley, T.M., 2010. Diversity and reductive evolution of mitochondria among microbial eukaryotes. *Philos. Trans. R. Soc. B-Biol. Sci.* 365, 713–727.
- Hopkins, J., Fowler, R., Krishna, S., Wilson, I., Mitchell, G., Bannister, L., 1999. The plastid in *Plasmodium falciparum* asexual blood stages: a three-dimensional ultrastructural analysis. *Protist* 150, 283–295.
- Howe, C.J., 1992. Plastid origin of an extrachromosomal DNA molecule from *Plasmodium*, the causative agent of malaria. *J. Theor. Biol.* 158, 199–205.
- Howe, C.J., Ellen, R., Nisbet, R., Barbrook, A.C., 2008. The remarkable chloroplast genome of dinoflagellates. *J. Exp. Bot* 59, 1035–1045.
- Jaffe, E.K., Shanmugam, D., Gardberg, A., Dieterich, S., Sankaran, B., Stewart, L.J., Myler, P.J., Roos, D.S., 2011. Crystal structure of *Toxoplasma gondii* porphobilinogen synthase: Insights on octameric structure and porphobilinogen formation. *J. Biol. Chem.* 286, 15298–15307.
- Janouskovec, J., Horak, A., Obornik, M., Lukes, J., Keeling, P.J., 2010. A common red algal origin of the apicomplexan, dinoflagellate, and heterokont plastids. *Proc. Natl Acad. Sci. U S A* 107, 10949–10954.
- Jomaa, H., Wiesner, J., Sanderbrand, S., Altincicek, B., Weidmeyer, C., Hintz, M., Turbachova, I., Eberl, M., Zeidler, J., Lichtenthaler, H.K., Soldati, D., Beck, E., 1999. Inhibitors of the nonmevalonate pathway of isoprenoid biosynthesis as antimalarial drugs. *Science* 285, 1573–1576.
- Joseph, S.J., Fernandez-Robledo, J.A., Gardner, M.J., El-Sayed, N.M., Kuo, C.H., Schott, E.J., Wang, H., Kissinger, J.C., Vasta, G.R., 2010. The Alveolate *Perkinsus marinus*: biological insights from EST gene discovery. *BMC Genomics* 11, 228.
- Kalanon, M., McFadden, G.I., 2010. Malaria, *Plasmodium falciparum* and its apicoplast. *Biochem. Soc. Trans.* 38, 775–782.
- Keeling, P.J., 2010. The endosymbiotic origin, diversification and fate of plastids. *Philos. Trans. R. Soc. B-Biol. Sci.* 365, 729–748.
- Kilejian, A., 1974. A unique histidine-rich polypeptide from the malaria parasite, *Plasmodium lophurae*. *J. Biol. Chem.* 249, 4650–4655.
- Kissinger, J.C., Kuo, C.-H., 2007. Evolution and comparative genomics of *Toxoplasma gondii*. In: Ajioka, J.W., Soldati, D. (Eds.), *Toxoplasma: Molecular and Cellular Biology*. Horizon Bioscience, Norfolk, UK, pp. 209–226.
- Köhler, S., Delwiche, C.F., Denny, P.W., Tilney, L.G., Webster, P., Wilson, R.J., Palmer, J.D., Roos, D.S., 1997. A plastid of probable green algal origin in Apicomplexan parasites. *Science* 275, 1485–1489.
- Kudryashev, M., Lepper, S., Stanway, R., Bohn, S., Baumeister, W., Cyrklaff, M., Frischknecht, F., 2010. Positioning of large organelles by a membrane-associated cytoskeleton in *Plasmodium* sporozoites. *Cell. Microbiol.* 12, 362–371.
- Kuvardina, O.N., Leander, B.S., Aleshin, V.V., Myl'nikov, A.P., Keeling, P.J., Simdyanov, T.G., 2002. The phylogeny of colpodellids (Alveolata) using small subunit rRNA gene sequences suggests they are the free-living sister group to apicomplexans. *J. Eukaryot. Microbiol.* 49, 498–504.
- La Peyre, J.F., Faisal, M., 1997. Development of a protein-free chemically defined culture medium for the propagation of the oyster pathogen *Perkinsus marinus*. *Parasite* 4, 67–73.
- Leander, B.S., Hoppenrath, M., 2008. Ultrastructure of a novel tube-forming, intracellular parasite of dinoflagellates: *Parvilicifera proocentri* sp. nov. (Alveolata, Myzozoa). *Eur J Protistol.* 44, 55–70.
- Leander, B.S., Keeling, P.J., 2003. Morphostasis in alveolate evolution. *Trends Ecol. Evol.* 18, 395–402.
- Lester, R.J.C., Davis, G.H.G., 1981. A new *Perkinsus* species (Apicomplexa, Perkinsea) from the Abalone *Haliotis ruber*. *J. Invertebr. Pathol.* 37, 181–187.
- Levine, N.D., 1978. *Perkinsus* gen.n. and other new taxa in the protozoan phylum Apicomplexa. *J. Parasitol.* 64, 549.
- Lim, L., Kalanon, M., McFadden, G.I., 2009. New proteins in the apicoplast membranes: time to rethink apicoplast protein targeting. *Trends Parasitol.* 25, 197–200.
- Lim, L., McFadden, G.I., 2010. The evolution, metabolism and functions of the apicoplast. *Philos. Trans. R. Soc. B-Biol. Sci.* 365, 749–763.
- Lin, Z., Fernández-Robledo, J.A., Cellier, M.F., Vasta, G.R., 2011. The natural resistance-associated macrophage protein from the protozoan parasite *Perkinsus marinus* mediates iron uptake. *Biochemistry* 50, 6340–6355.
- Lund, E.D., Chu, F.L., 2002. Phospholipid biosynthesis in the oyster protozoan parasite, *Perkinsus marinus*. *Mol. Biochem. Parasitol.* 121, 245–253.
- Lund, E.D., Chu, F.L., Soudant, P., Harvey, E., 2007. *Perkinsus marinus*, a protozoan parasite of the eastern oyster, has a requirement for dietary sterols. *Comp. Biochem. Physiol. A: Mol. Integr. Physiol.* 146, 141–147.
- Lund, E.D., Soudant, P., Chu, F.L., Harvey, E., Bolton, S., Flowers, A., 2005. Effects of Triclosan on growth, viability and fatty acid synthesis of the oyster protozoan parasite *Perkinsus marinus*. *Dis. Aquat. Organ.* 67, 217–224.
- Mackin, J.G., Owen, H.M., Collier, A., 1950. Preliminary note on the occurrence of a new protistan parasite, *Dermocystidium marinum* n. sp., in *Crassostrea virginica* (Gmelin). *Science* 111, 328–329.
- Mackin, J.G.R., Ray, S.M., 1966. The taxonomic relationships of *Dermocystidium marinum* Mackin, Owen and Collier. *J. Invertebr. Pathol.* 8, 544–545.
- Martens, C., Vandepoel, K., Van de Peer, Y., 2008. Whole-genome analysis reveals molecular innovations and evolutionary transitions in chromalveolate species. *Proc. Natl Acad. Sci. USA* 105, 3427–3432.
- Martin, W., Herrmann, R.G., 1998. Gene transfer from organelles to the nucleus: how much, what happens, and Why? *Plant Physiol.* 118, 9–17.
- Martin, W., Rujan, T., Richly, E., Hansen, A., Cornelsen, S., Lins, T., Leister, D., Stoebe, B., Hasegawa, M., Penny, D., 2002. Evolutionary analysis of *Arabidopsis*,

- cyanobacterial, and chloroplast genomes reveals plastid phylogeny and thousands of cyanobacterial genes in the nucleus. *Proc. Natl Acad. Sci. USA* 99, 12246–12251.
- Masuda, I., Matsuzaki, M., Kita, K., 2010. Extensive frameshift at all AGG and CCC codons in the mitochondrial cytochrome c oxidase subunit 1 gene of *Perkinsus marinus* (Alveolata; Dinoflagellata). *Nucleic Acids Res.* 38, 6186–6194.
- Matsuzaki, M., Kita, K., Nozaki, H., 2006. Orthologs of plastid isoprenoids biosynthesis pathway genes from *Perkinsus*. *J. Plant. Res.* 119 (suppl), 122.
- Matsuzaki, M., Kuroiwa, H., Kuroiwa, T., Kita, K., Nozaki, H., 2008. A cryptic algal group unveiled: a plastid biosynthesis pathway in the oyster parasite *Perkinsus marinus*. *Mol. Biol. Evol.* 25, 1167–1179.
- McFadden, G.I., Reith, M.E., Munholland, J., Lang-Unnasch, N., 1996. Plastid in human parasites. *Nature* 381, 482.
- McFadden, G.I., Roos, D.S., 1999. Apicomplexan plastids as drug targets. *Trends Microbiol.* 7, 328–333.
- McLaughlin, S.M., Tall, B.D., Shaheen, A., Elsayed, E.E., Faisal, M., 2000. Zoosporulation of a new *Perkinsus* species isolated from the gills of the softshell clam *Mya arenaria*. *Parasite* 7, 115–122.
- Miranda, K., Pace, D.A., Cintron, R., Rodrigues, J.C., Fang, J., Smith, A., Rohloff, P., Coelho, E., de Haas, F., de Souza, W., Coppens, I., Sibley, L.D., Moreno, S.N., 2010. Characterization of a novel organelle in *Toxoplasma gondii* with similar composition and function to the plant vacuole. *Mol. Microbiol.* 76, 1358–1375.
- Moore, R.B., Obornik, M., Janouskovec, J., Chrudimsky, T., Vancova, M., Green, D.H., Wright, S.W., Davies, N.W., Bolch, C.J., Heimann, K., Slapeta, J., Hoegh-Guldberg, O., Logsdon, J.M., Carter, D.A., 2008. A photosynthetic alveolate closely related to apicomplexan parasites. *Nature* 451, 959–963.
- Moss, J.A., Xiao, J., Dungan, C.F., Reece, K.S., 2008. Description of *Perkinsus beihaiensis* n. sp., a new *Perkinsus* sp. parasite in oysters of Southern China. *J. Eukaryot Microbiol.* 55, 117–130.
- Muller, J., Hemphill, A., 2011. Drug Target Identification in Intracellular and Extracellular Protozoan Parasites. *Curr Top Med Chem.* [Epub ahead of print].
- Murrell, A., Kleeman, S.N., Barker, S.C., Lester, R.J.G., 2002. Synonymy of *Perkinsus olseni* Lester & Davis, 1981 and *Perkinsus atlanticus* Azevedo, 1989 and an update on the phylogenetic position of the genus *Perkinsus*. *Bull. EAFP.* 22, 258–265.
- Nagamune, K., Hicks, L.M., Fux, B., Brossier, F., Chini, E.N., Sibley, L.D., 2008a. Abscisic acid controls calcium-dependent egress and development in *Toxoplasma gondii*. *Nature* 451, 207–210.
- Nagamune, K., Xiong, L., Chini, E., Sibley, L.D., 2008b. Plants, endosymbionts and parasites: Abscisic acid and calcium signaling. *Commun Integr Biol.* 1, 62–65.
- Nagaraj, V.A., Prasad, D., Arumugam, R., Rangarajan, P.N., Padmanaban, G., 2010. Characterization of coproporphyrinogen III oxidase in *Plasmodium falciparum* cytosol. *Parasitol Int.* 59, 121–127.
- Nambara, E., Marion-Poll, A., 2005. Abscisic acid biosynthesis and catabolism. *Annu. Rev. Plant Biol.* 56, 165–185.
- Nishi, M., Hu, K., Murray, J.M., Roos, D.S., 2008. Organellar dynamics during the cell cycle of *Toxoplasma gondii*. *J. Cell Sci.* 121, 1559–1568.
- Norén, F., Moestrup, O., Rehnstam-Holm, A.S., 1999. *Parvilucifera infectans* Norén et Moestrup gen. et sp. nov. (Perkinsozoa phylum nov.): A parasitic flagellate capable of killing toxic microalgae. *Eur J Protistol.* 35, 233–254.
- Pain, A., Renaud, H., Berriman, M., Murphy, L., Yeats, C.A., Weir, W., Kerhoun, A., Aslett, M., Bishop, R., Bouchier, C., Cochet, M., Coulson, R.M., Cronin, A., de Villiers, E.P., Fraser, A., Fosker, N., Gardner, M., Goble, A., Griffiths-Jones, S., Harris, D.E., Katzer, F., Larke, N., Lord, A., Maser, P., McKellar, S., Mooney, P., Morton, F., Nene, V., O'Neil, S., Price, C., Quail, M.A., Rabinovitsch, E., Rawlings, N.D., Rutter, S., Saunders, D., Seeger, K., Shah, T., Squares, R., Squares, S., Tivey, A., Walker, A.R., Woodward, J., Dobbelaere, D.A., Langsley, G., Rajandream, M.A., McKeever, D., Shiels, B., Tait, A., Barrell, B., Hall, N., 2005. Genome of the host-cell transforming parasite *Theileria annulata* compared with *T. parva*. *Science* 309, 131–133.
- Pecher, W.T., Alavi, M.R., Schott, E.J., Fernández-Robledo, J.A., Roth, L., Berg, S.T., Vasta, G.R., 2008. Assessment of the northern distribution range of selected *Perkinsus* species in eastern oysters (*Crassostrea virginica*) and hard clams (*Mercentaria mercenaria*) with the use of PCR-based detection assays. *J. Parasitol.* 94, 410–422.
- Perkins, F.O., 1976. Zoospores of the oyster pathogen, *Dermocystidium marinum*. I. Fine structure of the conoid and other sporozoan-like organelles. *J. Parasitol.* 62, 959–974.
- Perkins, F.O., 1996. The structure of *Perkinsus marinus* (Mackin, Owen & Collier, 1950) Levine, 1978 with comments on taxonomy and phylogeny of *Perkinsus* spp. *J. Shellfish Res.* 15, 67–87.
- Ralph, S.A., Foth, B.J., Hall, N., McFadden, G.I., 2004a. Evolutionary pressures on apicoplast transit peptides. *Mol. Biol. Evol.* 21, 2183–2194.
- Ralph, S.A., Van Dooren, G.G., Waller, R.F., Crawford, M.J., Fraunholz, M.J., Foth, B.J., Tonkin, C.J., Roos, D.S., McFadden, G.I., 2004b. Tropical infectious diseases: Metabolic maps and functions of the *Plasmodium falciparum* apicoplast. *Nat. Rev. Microbiol.* 2, 203–216.
- Reece, K.S., Bushek, D., Hudson, K.L., Graves, J.E., 2001. Genetic distribution of *Perkinsus marinus* genetic strains along the Atlantic and Gulf coasts of the USA. *Mar. Biol.* 139, 1047–1055.
- Reece, K.S., Siddall, M.E., Burreson, E.M., Graves, J.E., 1997. Phylogenetic analysis of *Perkinsus* based on actin gene sequences. *J. Parasitol.* 83, 417–423.
- Reggiori, F., Klionsky, D.J., 2005. Autophagosomes: biogenesis from scratch? *Curr. Opin. Cell Biol.* 17, 415–422.
- Robledo, J.A., Nunes, P.A., Cancela, M.L., Vasta, G.R., 2002. Development of an *in vitro* clonal culture and characterization of the rDNA locus of *Perkinsus atlanticus*, a protistan parasite of the clam *Tapes decussatus*. *J. Eukaryot. Microbiol.* 49, 414–422.
- Robledo, J.A.F., Wright, A.C., Marsh, A.G., Vasta, G.R., 1999. Nucleotide sequence variability in the nontranscribed spacer of the rRNA locus in the oyster parasite *Perkinsus marinus*. *J. Parasitol.* 85, 650–656.
- Saldarriaga, J.F., McEan, M.L., Fast, N.M., Taylor, F.J.R., Keeling, P.J., 2003. Multiple protein phylogenies show that *Oxyrrhis marina* and *Perkinsus marinus* are early branches of the dinoflagellate lineage. *Int. J. Syst. Evol. Microbiol.* 53, 355–365.
- Saldarriaga, J.F., Taylor, F.J.R.M., Cavalier-Smith, T., Menden-Deuer, S., Keeling, P.J., 2004. Molecular data and the evolutionary history of dinoflagellates. *Europ J Protol.* 40, 85–111.
- Sánchez-Puerta, M.V., Lippmeier, J.C., Apt, K.E., Delwiche, C.F., 2007. Plastid genes in a non-photosynthetic dinoflagellate. *Protist.* 158, 105–117.
- Schott, E.J., Vasta, G.R., 2003. The PmSOD1 gene of the protistan parasite *Perkinsus marinus* complements the sod2Delta mutant of *Saccharomyces cerevisiae*, and directs an iron superoxide dismutase to mitochondria. *Mol. Biochem. Parasitol.* 126, 81–92.
- Schott, E.J., Pecher, W.T., Robledo, J.A.F., Vasta, G.R., 2004. Ascorbate peroxidase and the antioxidant repertoire of *Perkinsus marinus*. In: XIV Molecular Parasitology Meeting, Woods Hole, MA, USA, p. 268A.
- Schwartz, S.H., Qin, X., Zeevaert, J.A., 2003. Elucidation of the indirect pathway of abscisic acid biosynthesis by mutants, genes, and enzymes. *Plant Physiol.* 131, 1591–1601.
- Seeber, F., Soldati-Favre, D., 2010. Metabolic pathways in the apicoplast of apicomplexa. *Int Rev Cell Mol Biol.* 281, 161–228.
- Sheiner, L., Soldati-Favre, D., 2008. Protein trafficking inside *Toxoplasma gondii*. *Traffic.* 9, 636–646.
- Siddall, M.E., Reece, K.S., Graves, J.E., Burreson, E.M., 1997. 'Total evidence' refutes the inclusion of *Perkinsus* species in the phylum Apicomplexa. *Parasitology* 115, 165–176.
- Slamovits, C.H., Keeling, P.J., 2008. Plastid-derived genes in the nonphotosynthetic alveolate *Oxyrrhis marina*. *Mol. Biol. Evol.* 25, 1297–1306.
- Soldati, D., 1999. The apicoplast as a potential therapeutic target in and other apicomplexan parasites. *Parasitol Today.* 15, 5–7.
- Spork, S., Hiss, J.A., Mandel, K., Sommer, M., Kooij, T.W., Chu, T., Schneider, G., Maier, U.G., Przyborski, J.M., 2009. An unusual ERAD-like complex is targeted to the apicoplast of *Plasmodium falciparum*. *Eukaryot Cell.* 8, 1134–1145.
- Stelzer, K., El-Sayed, N.M., Seeber, F., 2007. The expression of a plant-type ferredoxin redox system provides molecular evidence for a plastid in the early dinoflagellate *Perkinsus marinus*. *Protist.* 158, 119–130.
- Sunila, I., Hamilton, R.M., Dungan, C.F., 2001. Ultrastructural characteristics of the *in vitro* cell cycle of the protozoan pathogen of oysters, *Perkinsus marinus*. *J. Eukaryot. Microbiol.* 48, 348–361.
- Tasumi, S., Vasta, G.R., 2007. A galectin of unique domain organization from hemocytes of the eastern oyster (*Crassostrea virginica*) is a receptor for the protistan parasite *Perkinsus marinus*. *J. Immunol.* 179, 3086–3098.
- Teles-Grilo, M.L., Duarte, S.M., Tato-Costa, J., Gaspar-Maia, A., Oliveira, C., Rocha, A.A., Marques, A., Cordeiro-da-Silva, A., Azevedo, C., 2007a. Molecular karyotype analysis of *Perkinsus atlanticus* (Phylum Perkinsozoa) by pulsed field gel electrophoresis. *Eur J Protistol.* 43, 315–318.
- Teles-Grilo, M.L., Tato-Costa, J., Duarte, S.M., Maia, A., Casal, G., Azevedo, C., 2007b. Is there a plastid in *Perkinsus atlanticus* (Phylum Perkinsozoa)? *Eur J Protistol.* 43, 163–167.
- Thompson, P.C., Rosenthal, B.M., Hare, M.P., 2011. An evolutionary legacy of sex and clonal reproduction in the protistan oyster parasite *Perkinsus marinus*. *Infect Genet Evol.* 11, 598–609.
- Tonhosolo, R., D'Alexandri, F.L., de Rosso, V.V., Gazarini, M.L., Matsumura, M.Y., Peres, V.J., Merino, E.F., Carlton, J.M., Wunderlich, G., Mercadante, A.Z., Kimura, E.A., Katzin, A.M., 2009. Carotenoid biosynthesis in intraerythrocytic stages of *Plasmodium falciparum*. *J. Biol. Chem.* 284, 9974–9985.
- Tonkin, C.J., Foth, B.J., Ralph, S.A., Struck, N., Cowman, A.F., McFadden, G.I., 2008a. Evolution of malaria parasite plastid targeting sequences. *Proc. Natl Acad. Sci. USA* 105, 4781–4785.
- Tonkin, C.J., Kalanon, M., McFadden, G.I., 2008b. Protein targeting to the malaria parasite plastid. *Traffic.* 9, 166–175.
- Tonkin, C.J., Pearce, J.A., McFadden, G.I., Cowman, A.F., 2006a. Protein targeting to destinations of the secretory pathway in the malaria parasite *Plasmodium falciparum*. *Curr. Opin. Microbiol.* 9, 381–387.
- Tonkin, C.J., Roos, D.S., McFadden, G.I., 2006b. N-terminal positively charged amino acids, but not their exact position, are important for apicoplast transit peptide fidelity in *Toxoplasma gondii*. *Mol. Biochem. Parasitol.* 150, 192–200.
- Vaishnav, S., Striepen, B., 2006. The cell biology of secondary endosymbiosis—how parasites build, divide and segregate the apicoplast. *Mol. Microbiol.* 61, 1380–1387.
- van der Giezen, M., Tovar, J., Clark, C.G., 2005. Mitochondrion-derived organelles in protists and fungi. *Int. Rev. Cytol.* 244, 175–225.
- Villalba, A., Reece, K.S., Ordás, A., Casas, S.M., Figueroa, A., 2004. Perkinsosis in molluscs: a review. *Aquat. Living Res.* 17, 411–432.
- Waller, R.F., Keeling, P.J., Donald, R.G., Striepen, B., Handman, E., Lang-Unnasch, N., Cowman, A.F., Besra, G.S., Roos, D.S., McFadden, G.I., 1998. Nuclear-encoded proteins target to the plastid in *Toxoplasma gondii* and *Plasmodium falciparum*. *Proc Natl Acad Sci USA* 95, 12352–12357.
- Waller, R.F., Slamovits, C.H., Keeling, P.J., 2006. Lateral gene transfer of a multigene region from cyanobacteria to dinoflagellates resulting in a novel plastid-targeted fusion protein. *Mol. Biol. Evol.* 23, 1437–1443.

- Weiss, L.M., Kim, K., 2007. *Toxoplasma gondii*: the model apicomplexan: perspectives and methods. Elsevier, San Diego, CA, USA.
- Wiesner, J., Reichenberg, A., Heinrich, S., Schlitzer, M., Jomaa, H., 2008. The plastid-like organelle of apicomplexan parasites as drug target. *Curr. Pharm. Des.* 14, 855–871.
- Williams, B.A., Keeling, P.J., 2003. Cryptic organelles in parasitic protists and fungi. *Adv. Parasitol.* 54, 9–68.
- Williamson, D.H., Gardner, M.J., Preiser, P., Moore, D.J., Rangachari, K., Wilson, R.J., 1994. The evolutionary origin of the 35 kb circular DNA of *Plasmodium falciparum*: new evidence supports a possible rhodophyte ancestry. *Mol. Gen. Genet.* 243, 249–252.
- Wilson, R.J., Denny, P.W., Preiser, P.R., Rangachari, K., Roberts, K., Roy, A., Whyte, A., Strath, M., Moore, D.J., Moore, P.W., Williamson, D.H., 1996. Complete gene map of the plastid-like DNA of the malaria parasite *Plasmodium falciparum*. *J. Mol. Biol.* 261, 155–172.
- Wright, A., Ahmed, H., Gauthier, J., Silva, A., Vasta, G., 2002. CDNA cloning and characterization of two iron superoxide dismutases from the oyster parasite *Perkinsus marinus*. *Mol. Biochem. Parasitol.* 123, 73–77.
- Xu, P., Widmer, G., Wang, Y., Ozaki, L.S., Alves, J.M., Serrano, M.G., Puiu, D., Manque, P., Akiyoshi, D., Mackey, A.J., Pearson, W.R., Dear, P.H., Bankier, A.T., Peterson, D.L., Abrahamsen, M.S., Kapur, V., Tzipori, S., Buck, G.A., 2004. The genome of *Cryptosporidium hominis*. *Nature* 431, 1107–1112.
- Xu, X.M., Moller, S.G., 2011. Iron–sulfur clusters: Biogenesis, molecular mechanisms, and their functional significance. *Antioxid. Redox Signal.* 15, 271–307.
- Yorimitsu, T., Klionsky, D.J., 2005. Autophagy: molecular machinery for self-eating. *Cell Death Differ.* 12, 1542–1552.
- Yu, M., Kumar, T.R., Nkrumah, L.J., Coppi, A., Retzlaff, S., Li, C.D., Kelly, B.J., Moura, P.A., Lakshmanan, V., Freundlich, J.S., Valderramos, J.C., Vilcheze, C., Siedner, M., Tsai, J.H., Falkard, B., Sidhu, A.B., Purcell, L.A., Gratraud, P., Kremer, L., Waters, A.P., Schiehsler, G., Jacobus, D.P., Janse, C.J., Ager, A., Jacobs Jr., W.R., Sacchettini, J.C., Heussler, V., Sinnis, P., Fidock, D.A., 2008. The fatty acid biosynthesis enzyme FabI plays a key role in the development of liver-stage malarial parasites. *Cell Host Microbe* 4, 567–578.
- Zhang, H., Dungan, C.F., Lin, S., 2011. Introns, alternative splicing, spliced leader trans-splicing and differential expression of pcna and cyclin in *Perkinsus marinus*. *Protist.* 162, 154–167.
- Zhang, H., Hou, Y., Miranda, L., Campbell, D.A., Sturm, N.R., Gaasterland, T., Lin, S., 2007. Spliced leader RNA trans-splicing in dinoflagellates. *Proc. Natl Acad. Sci. USA* 104, 4618–4623.
- Zhu, G., Marchewka, M.J., Keithly, J.S., 2000. *Cryptosporidium parvum* appears to lack a plastid genome. *Microbiology* 146, 315–321.
- Zuegge, J., Ralph, S., Schmuker, M., McFadden, G.I., Schneider, G., 2001. Deciphering apicoplast targeting signals – feature extraction from nuclear-encoded precursors of *Plasmodium falciparum* apicoplast proteins. *Gene* 280, 19–26.